

A
07/322604

- 1 -

MICROORGANISMS AND PLASMIDS FOR
2,4-DICHLOROPHENOXYACETIC ACID (2,4-D)
MONOOXYGENASE FORMATION AND PROCESS FOR
THE PRODUCTION OF THESE PLASMIDS AND STRAINS

5 This is a continuation-in-part of PCT Application
No. PCT/DE87/00392, filed August 28, 1987 ~~(Attorney's~~
~~Docket SCH 920)~~.

Background of the Invention

10 The present invention relates to the production,
by genetic engineering, of plasmids and bacterial strains
containing the gene *tfdA*, or a gene essentially identical
to *tfdA*, on a short, exactly characterizable DNA segment.
The novel plasmids and microorganisms are especially well
suited for the production of 2,4-D-monooxygenase, as well as
15 for serving as starting compounds for the genetically
engineered transfer of the 2,4-D-degrading properties of
this enzyme to various organisms (including the thus-
attainable 2,4-D-tolerance in genetically transformed
plants).

20 2,4-D-Monooxygenase is an enzyme catalyzing in
many 2,4-D-degrading organisms the first step in the
metabolizing of 2,4-dichlorophenoxyacetic acid (2,4-D).
Among the 2,4-D-degrading organisms belong, in particular,
soil bacteria, such as, for example, *Acinetobacter*,
25 *Alcaligenes*, *Arthrobacter*, *Cyrenebacterium* and *Pseudomonas*
[compare, in this connection, G.R. Bell, Can. J. Microbiol.
3 : 821 (1957); J.M. Bollag et al., J. Agric. Food Chem.
16 : 826 (1968); R.H. Don et al., J. Bacteriol. 145 : 681
(1981); W.C. Evans et al., Proc. Biochem. Soc. Biochem. J.
30 57 : 4 (1954); W.C. Evans et al., Biochem. J. 122: 543
(1971); R.P. Fisher et al., J. Bacteriol. 135 : 798

(1978); T.I. Steenson et al., J. Gen. Microbiol. 16 : 146
(1957); J.M. Tiedje et al., J. Agric. Food Chem. 17 : 1080
(1969); J.E. Tyler et al., Appl. Microbiol. 28 : 181 (1974);
J.M. Tiedje et al., J. Agric. Food Chem. 17 : 1021
5 (1969)]. They play a significant part in the detoxifica-
tion of the soil and of the wastewaters from halogenated
aromatic compounds as they occur especially among the
agriculturally utilized pesticides and herbicides.

In the group of the 2,4-D-degrading wild-type
10 bacteria, the strain *Alcaligenes eutrophus* JMP134 is the
most well-known and best-characterized. This strain
harbors the plasmid pJP4 having a size of about 80 kilo-
bases and containing all of the genes important for
degradation of 2,4-D (R.H. Don et al., loc. cit.). The
15 plasmid pJP4 has also been isolated and characterized
recently [R.H. Don et al., J. Bacteriol. 161 : 466 (1985)].

Five genes participating in 2,4-D degradation,
denoted as *tfdB*, *tfdC*, *tfdD*, *tfdE* and *tfdF*, were localized
by transposon mutagenesis and cloned in *E. coli*. R.H. Don
20 et al. [J. Bacteriol. 161 : 85 (1985)] succeeded in attrib-
uting enzyme function to four genes by biochemical studies.
However, thus far attempts have been unsuccessful regarding
localizing of gene *tfdA* on the plasmid pJP4 or on a sub-
fragment of the latter, and to isolate this gene and to
25 clarify its structure.

Summary of the Invention

It has now been possible for the first time thanks
to the present invention to isolate, to clone, and to char-
acterize the gene *tfdA*. It has thus become possible to
30 make this gene available for transfer to other organisms
with the objective of expressing the *tfdA*-encoded 2,4-D-
monooxygenase in these organisms. The latter includes micro-
organisms, as well as higher organisms, such as, for example,
plants.

35 The controlled transfer of *tfdA* to other micro-
organisms offers the possibility of broadening the spectrum
of compounds degradable by these organisms. As far as
bacteria are concerned, transfer and expression of genes
of the entire 2,4-D degradation has been disclosed in

3

/

J. Bacteriol. 145 : 681 (1981) and in Arch. Microbiol. 134 : 92 (1983). However, no attempt has as yet been made to express an isolated *tfdA* gene in gram-negative bacteria, such as *Pseudomonas* or *Alcaligenes*.

5 No known attempt has been made in connection with plants, either. Metabolism of 2,4-D by several types of plants has been reported by various sources [Weed Science 24 : 557 (1976) and Z. Pflanzenphysiol. 110⁴: 395 (1983)]. In lower concentrations, this compound
10 acts as an auxin-analog plant hormone and therefore is utilized in cell culturing technique; at higher concentrations, it acts on the plant cell as well as on the entire plant in the manner of a growth inhibitor, which is the reason why it is utilized as a herbicide. In a
15 haploid cell suspension culture of *Nicotiana glauca*, adaptation to rising concentrations of 2,4-D brought about tolerance with respect to this synthetic growth compound based on an increased rate of metabolism [M.H. Zenk, 1974, Haploids in Physiological and Biochemical Research, in Haploids in Higher Plants. Advances and Potential; Proceedings of the First International Symposium, Guelph, Ontario, Canada, p. 339].

B The *tfdA* gene codes for ~~2,4-D-monoamine oxygenase~~ ^{2,4-D-mono-oxygenase}, a polypeptide having the biological activity of bringing about the cleavage of the side chain of
25 2,4-D; therefore, the ability to inactivate 2,4-D can be transferred, with the aid of the gene obtained from bacteria, to all those plants lending themselves to such a manipulation by genetic engineering. Methods for transferring foreign genes to plants and their progeny are already known for several types of plants (M. De Block,
30 L. Herrera-Estrella, M. Van Montagu, J. Schell, and P. Zambryski, Expression of Foreign Genes in Regenerated Plants and in Their Progeny, EMBO J. 3 : 1681; as well as R.D. Shillito, M.W. Saul, J. Paszkowski, M. Muller, and J. Potrykus, 1985, High Efficiency Direct Gene Transfer to Plants, Bio/technology 3 : 1099).

4

Great hopes are harbored for the genetic transformation of plants, especially regarding the production of novel useful plants important to man [J.L. Marx, 1985, Plant Gene Transfer Becomes a Fertile Field, Science 230 : 1148]. In this context, the gene *tfdA* provides the availability of a novel, genetically transferable and in vivo selectable property in the form of tolerance against a growth inhibitor, as has been known heretofore merely for a few tolerances against antibiotics and, respectively, herbicides [R.T. Fraley et al., 1983, Expression of Bacterial Genes in Plant Cells, Proc. Natl. Acad. Sci. USA 80 : 4803; and Nature 317 : 741 (1985)].

The present invention, thus provides a heretofore unknown mutant using a conventional mutation method, from a bacterial strain capable of utilizing 2,4-D as the growth substrate, such as, for example, *Alcaligenes eutrophus* JMP134, the structural gene for ^{2,4-D monooxygenase}~~2,4-D monooxygenase~~ being inactivated in this mutant. This mutant is used, employing the known methods of DNA recombination in vitro, of transformation with recombinant DNA, and of conjugative transfer of DNA, for the selection of recombinant DNA containing *tfdA* or genes substantially identical to *tfdA*.

Another aspect of this invention relates to plasmids containing the gene *tfdA* or genes substantially identical to *tfdA* (e.g., hybridizing with the gene of this invention and coding for a protein which has the biological activity of the protein encoded by *tfdA*, e.g., its 2,4-D-monooxygenase activity), as well as plasmids containing parts of these genes including the promoter region.

A third aspect of this invention is to provide novel bacterial host/plasmid strains which can be used to produce 2,4-D-monooxygenase, or which can be used to transfer the 2,4-D-monooxygenase gene to other organisms.

A fourth aspect of this invention is the creation of transgenic organisms, preferably microorganisms and plants, which are resistant to the growth inhibiting effects of certain substituted or unsubstituted phenoxyacetic acids.

5 The plasmids of this invention can be produced by digesting DNA from wild-type bacteria exhibiting genes for metabolizing 2,4-D or compounds similar to 2,4-D with restriction endonucleases and coupling the thus-obtained DNA fragments, with the aid of a DNA ligase,
10 with a plasmid vector which latter has previously been converted into its linear form by the same restriction enzyme.

 Among the thus-formed recombinant plasmids, the tfdA-containing plasmids according to this invention can be
15 identified by introducing the plasmids into bacterial strains from which then the tfdA-containing clones can be directly selected based on a tfdA-conveyed ability.

 Such bacterial strains have the property of being able to exploit as sources of carbon and energy the products
20 formed in an enzymatic reaction in vivo by 2,4-D-monooxygenase, but not the substrates thereof.

 Examples of strains having the aforementioned ability are the tfdA mutants according to this invention wherein all genes are active for degradation of 2,4-
25 dichlorophenol, furthermore the strain *Alcaligenes eutrophus* JMP222 which exhibits degradation pathways for phenol, as well as *Pseudomonas* sp. B13 which can utilize 4-chlorophenol. However, a series of further bacterial strains which can degrade differently substituted phenols
30 and are encountered primarily among the group of gram-negative bacteria can conceivably be recipients for the selection and identification of cloned tfdA genes.

 Preferred cloning vectors are plasmids having a wide host range, capable of replicative growth even in
35 bacterial strains other than *E. coli*. However, plasmids capable of multiplying their DNA or parts of their DNA by integration into the genome of the host cell can likewise be used for this purpose.

6

A suitable process for introducing DNA into the living cells of the aforementioned bacterial strains is, first of all, the direct transformation of these strains insofar as methods therefor are known. Thus, transformation methods exist, for example, for *Pseudomonas* and related gram-negative bacteria [A.A. Mercer and J.S. Loutit, 1979, Transformation and Transfection of *Pseudomonas aeruginosa*: Effects of Metal Ions, J. Bacteriol. 140 : 37-42; A.M. Chakrabarty, J.R. Mylroie, D.A. Friello, and J.G. Vacca, 1975, Transformation of *Pseudomonas putida* and *Escherichia coli* with Plasmid-Linked Drug-Resistance Factor DNA, Proc. Natl. Acad. Sci. USA 72 : 3647 - 3651]. In contrast thereto, a procedure which is applicable to all gram-negative bacteria and is substantially more effective is the transformation of an *E. coli* strain according to a method known from the literature with subsequent conjugative transfer of the plasmids into the respective bacterial strains.

Mobilizable plasmid vectors are used with preference for the conjugative transfer of cloned DNA. The genes required for transfer are made available in this case either by so-called helper plasmids or by mobilizing strains specifically designed for this purpose.

The plasmids containing *tfdA* can be isolated according to known methods as unequivocally definable chemical compounds from the aforesaid bacterial strains obtained directly by selection on suitable growth substrates or from clones of *E. coli* identified by the aforementioned method or by other conventional testing systems as regards expression of *tfdA*.

The advantage of direct selection of *tfdA*-containing strains by a growth test resides, above all, in that it permits, in contrast to the previously known methods all of which are based on relatively expensive enzyme tests, the identification of a *tfdA*-containing plasmid among a very large number of various plasmids as they are produced with preference in the preparation of gene banks, i.e. the randomized cloning of DNA fragments of genomic DNA. The advantage thus resides in broad

7

applicability of the process, for cloning of *tfdA* genes is thus no longer restricted to wild-type strains carrying the gene *tfdA* on a readily isolable plasmid, but rather is made possible from almost all wild-type strains, even those containing the gene on a poorly accessible plasmid or on the chromosome.

The plasmids of this invention are obtained, for example, by cutting the plasmid pJP4, stemming from *Alcaligenes eutrophus* JMP134, on which the genes lie for degrading 2,4-D, with the restriction endonuclease HindIII, separating the thus-formed DNA fragments by electrophoresis, and linking the individual isolated fragments with the HindIII-cut and dephosphorylated vector pVK101. The mobilizing strain *E. coli* S17-1 is transformed with the recombinant DNA, and plasmid-containing strains are selected on a tetracycline-containing medium. From strains containing recombinant plasmids identified by restriction analysis, the plasmid DNA is transferred by conjugation to the above-mentioned *tfdA* mutant JMP134:Tn5-2, and expression of the cloned *tfdA* gene is confirmed by growth of the host strain on 2,4-D-containing minimal medium. In this way, the plasmid pVJH21 can be identified containing the gene *tfdA* on a HindIII fragment having a size of 21 kilobases, stemming from pJP4. The identity of the cloned fragment can be confirmed by isolation of the plasmid and subsequent restriction analysis.

The plasmids according to this invention can furthermore be prepared by subcloning from recombinant plasmids containing the gene *tfdA*. For this purpose, these plasmids are digested with one or several restriction endonucleases and the thus-formed fragments are linked with the aid of a DNA ligase with a vector plasmid that has been converted into its linear form with the same restriction endonucleases. Plasmids that contain *tfdA* can be selected from the thus-formed plasmids in the way described above.

Thus, for example, the plasmid pGJS3 can be produced by linking DNA fragments of the plasmid pVJH21, formed with SacI, with the SacI-cut vector plasmid pGSS33. Among the number of newly combined plasmids, those can
5 be selected which contain an intact tfdA gene by transferring the recombinant plasmid DNA first by transformation into the mobilizing strain E. coli S17-1 and then from there by conjugation into the tfdA mutant JMP134:Tn5-2. By selection on 2,4-D-containing medium, plasmids are
10 identified which contain a SacI insert having a size of 3 kilobases, the origin of which can be clearly traced back by restriction analysis to the HindIII fragment, having a size of 21 kilobases, from pVJH21, and thus to pJP4.

By subcloning DNA segments containing tfdA,
15 those plasmids according to the invention can also be produced which differ in the type of plasmid vectors utilized, in dependence on the pursued target of usage.

Thus, for example, the 3-kilobase SacI fragment from pGJS3 can be transcloned into the vector pKT231,
20 thus forming the plasmids pKJS31 and pKJS32, offering as contrasted to pGJS3 advantages for the further characterization of tfdA due to a more favorable restriction map and the kanamycin resistance well expressed in many gram-negative bacteria. A preferred method for
25 confirming tfdA expression by pKT231-derived plasmids as described hereinbelow is the conjugative transfer from E. coli S17-1 into the strain *Alcaligenes eutrophus* JMP222 with subsequent testing for utilization of phenoxyacetic acid as the growth substrate. This system
30 has the advantages over the tfdA mutants that in this recipient strain the conjugative transfer takes place at higher efficiency and, in the kanamycin resistance, a further selectable marker is made available.

By subcloning, tfdA-containing DNA fragments can also be incorporated into expression vectors on which tfdA genes are expressed with the aid of a foreign (heterologous) promoter.

5 Thus, it is possible, for example, to incorporate
the 2.8 kilobase sized SacI/SalI fragment, the 2.0 kilobase
sized BamHI/SalI fragment, and the 1.4 kilobase sized
10 XbaI/SalI fragment into the expression vectors pT7-5 and
pT7-6 directly adjoining a phage promoter by means of which
gene expression can be activated in a controlled fashion
with the aid of a promoter-specific RNA polymerase.
The tfdA gene product expressed by the recombinant
plasmids pTJSS'035, pTJS'B435 and pTJS'X535, the pro-
duction of which will be described in Examples 8, 9 and 10,
15 can be identified by specific labeling with radioactive
methionine and subsequent gel electrophoresis. The
system can furthermore be used for producing, in an E. coli
strain, large amounts of 2,4-D-monooxygenase, which substan-
tially facilitates purification and subsequent protein-
20 chemical and enzymatic characterization of the gene
product as compared with its isolation from the wild-type
strain whereby additional biotechnological usage possibili-
ties become accessible to research.

Subcloning of tfdA-containing DNA fragments in
25 phage vectors of the type of the M13 phage makes it
possible to produce single-stranded DNA. The latter can
be utilized for determining the base sequence according to
the method of Sanger [F. Sanger, S. Nicklen, and A.R.
Coulson, 1977, DNA Sequencing with Chain-Terminating
30 Inhibitors, Proc. Natl. Acad. Sci. USA 74 : 5463]; it can
furthermore serve for the controlled mutagenesis of
individual bases by means of oligonucleotides, a conven-
tional method permitting the production of restriction
scission sites in a gene or the alteration of the amino
35 acid sequence. The insertion of restriction sites in genes

broadens the possibilities of application of the latter because cutting sites are thereby created for the incorporation of foreign promoters and the linkage of gene fragments for the formation of fusion proteins, as they
5 are required, for example, for the expression of prokaryotic genes in eukaryotes.

Subcloning of the 2.8-kilobase SacI/SalI fragment from pKJS32 into the phage vectors M13tg130 and M13tg131, as described in Example 16, represents a pos-
10 sibility for making the gene tfdA accessible to sequencing. By modification of the double-stranded DNA of the recombinant phages MJSS'030 and MJSS'031, additional alterations can be performed on the insert DNA.

The plasmids according to this invention can
15 furthermore evolve by modification from tfdA-containing plasmids. By treating such plasmids with one or several restriction endonucleases, optional treatment with exonucleases, and subsequent reassembly of the DNA ends to an annular molecule, deleted plasmids are obtained, i.e.
20 plasmids reduced in size by a specific DNA segment, which can contain a furthermore intact tfdA gene.

Thus, it is possible, for example, by the removal of defined DNA fragments from the plasmids pKJS31 and pKJS32, to shorten the insert contained therein,
25 having a size of 3 kilobases, on one side, up to the XbaI scission site and, on the other side, up to the SalI scission site, without losing the activity of tfdA during this step. The thus-produced plasmids pKJSB330, pKJS(X)630 and pKJS32RHAS', described in Examples 4, 5
30 and 7, are capable of expressing tfdA gene activity. In this way, the position of the gene can be delimited to an XbaI/SalI fragment of the size of 1.4 kilobases.

11

By the deletion, disclosed in Example 16, of variously long DNA segments from the double-stranded forms of the recombinant phages MJSS'030 and MJSS'031 by means of exonuclease III, a number of phages can be produced wherein respectively different regions of the cloned DNA lie closest to the starting point for sequencing. The sequence of a 2-kilobase BamHI/SalI fragment can then be composed of the partial regions sequenced in overlapping relationship. From a knowledge of the base sequence, further properties of the tfdA-containing DNA can then be derived, such as, for example, the position and length of the encoding region and the position of restriction scission sites, which can be of considerable utility for the further use of tfdA.

By deletion of DNA fragments from tfdA-containing plasmids and, respectively, by subcloning DNA fragments, it is also possible to obtain further plasmids according to this invention which then merely contain parts of a tfdA gene. Thus, for example, by subcloning the 1.5-kilobase EcoRI/BamHI fragment from pKJS32 into pKT231, the plasmid pKJEAB130 is obtained which merely contains the 5' end of tfdA, i.e. the promoter and the sequence coding for the N-terminus, and which is no longer capable of expressing an enzymatically active gene product. Parts of genes can be utilized for the construction of plasmids wherein the 5' end of a coding DNA, including its promoter, are linked in the same reading raster with the 3' end of another coding DNA and thus lead to expression of a so-called fusion protein in those organisms recognizing the respective promoter.

Another possibility of modifying the plasmids of this invention resides in the insertion of a foreign DNA segment. The insertion can serve for the introduction of new functional DNA sequences, e.g. of restriction scission sites, promoters, resistance genes, translation

and transcription stop signals. The insertion of a portion of a foreign gene sequence can also lead to the formation of fusion proteins.

5 One example of insertion of a foreign DNA is the insertion of the omega fragment into the BglII scission site of the plasmid pTJS'X535 located in the center of the coding region of tfdA (Example 11). With this fragment, translation stop codons are incorporated in all reading rasters and transcription stop signals
10 are incorporated together with a selectable resistance to antibiotics. As can be confirmed by specific radioactive labeling of the gene product as well as by in vivo enzyme test for 2,4-D-monooxygenase (Example 15), this results in the expression of a shortened, no longer
15 enzymatically active protein by the recombinant plasmid pTJS'X535omega.

Cloned fragments containing tfdA or parts of tfdA can serve for the detection of homologous DNA sequences and thus for finding tfdA genes in other
20 organisms. For this purpose, these fragments are excised from the plasmids of this invention with the aid of suitable restriction enzymes, isolated, and labeled in accordance with methods known from the literature, for example by incorporation of radionuclides. By means
25 of the conventional hybridization method, fragments can be identified in the entire DNA of an organism or in a gene bank produced therefrom, which fragments exhibit homology with tfdA. In this way, it is possible to recognize even among a population of various organisms
30 those which contain such a tfdA-homologous sequence. The process can be utilized for tracking down novel 2,4-D-degrading organisms as well as for the discovery of tfdA genes in these organisms.

It is possible to transfer *tfdA* genes to other organisms either directly with the aid of the inventive plasmids insofar as they are suitable for this purpose, or after modification by means of conventional genetic engineering techniques. The transfer can pursue the objective of expressing the *tfdA*-coded 2,4-D-monooxygenase in these organisms and thus to imbue the latter with the property of being able to degrade 2,4-D or compounds similar to 2,4-D.

Thus it is possible, for example, by means of the conjugative transfer of *tfdA*-containing plasmids with a wide host range into various gram-negative bacteria, described in Example 14, to impart to these bacteria the property of reacting 2,4-D or compounds similar to 2,4-D to the corresponding phenol. This, in conjunction with a degradation activity inherent in the respective strain, can lead to a broadening of the spectrum of compounds degradable by this strain. The strain *Alcaligenes eutrophus* JMP222 exhibits, for example, genes for metabolizing phenol. The expression of *tfdA* thus conveys to this strain the entirely novel possibility of utilizing phenoxyacetic acid as the growth substrate. The strain *Pseudomonas* B13 can metabolize phenol as well as 4-chlorophenol, and receives, by *tfdA*, in an analogous fashion the capability of growing on phenoxyacetic acid and 4-chlorophenoxyacetic acid. Likewise possible is the degradation of variously substituted phenoxyacetic acids, such as 4-chloro-2-methylphenoxyacetic acid, 2,4,5-trichlorophenoxyacetic acid, and similar compounds, by organisms containing *tfdA* and capable of reacting the corresponding phenols. Even though the thereby attainable broadening of the spectrum of degradation activities of an organism will not in every case lead to the creation of a novel property heretofore not realized in nature, it can yet serve for qualitative improvement insofar as strains can

thereby be constructed having a degrading power that is higher as compared with wild-type strains, or strains having higher tolerance against possibly toxic substrates or products of a degradation pathway. Such strains would
5 prove useful specifically under conditions prevailing in artificial systems, rather than in a natural environment, such as, for example, in clarification systems for the purification of industrial wastewaters. An overview has recently been published regarding the yields attainable
10 heretofore in the field of microbial degradation of halogenated aromatic compounds with the aid of newly designed strains [D. Ghosal, I.-S. You, D.K. Chatterjee, A.M. Chakrabarty, 1985, Microbial Degradation of Halogenated Compounds; Science 228 : 135-142].

15 A further possibility of utilizing the gene *tfdA* resides in its transfer to plants. Methods for the transformation of plants with foreign DNA are known in the literature. Thus far, there are two primary methods which can be used for transferring exogenous genes to plants: One method uses the
20 functions of the Ti plasmid from *Agrobacterium tumefaciens*; the other is based on the transformation of vegetable protoplasts with the aid of physical and chemical means and agents, such as "electroporation", thermal shock, treatment with calcium chloride or polyethylene glycol.

25 While transformation of plant cells by either method is always possible, in principle, the regeneration of entire plants from individual cells or cell tissues is not achieved in every instance. At present, this represents the primary obstacle to the manipulation of monocotyledonous
30 plants by genetic engineering.

A preferred objective of plant transformation with *tfdA* is the expression of the gene characteristic, i.e. the 2,4-D-monooxygenase activity, in the transformed plant, thus enabling the plant to degrade the derivatives
35 of phenoxyacetic acid, acting as an auxin-analogous plant

hormone at low concentrations and as a growth inhibitor at high concentrations, to the phytochemically ineffective phenols. Accordingly, this property would have significance as a selectable marker for genetically transformed plants.

5 In order for a prokaryotic gene, such as *tfdA*, to be expressed in plant cells, it must be coupled with the plant-specific signal sequences necessary for trans-
cription and translation. See Example 17. It has been mentioned
10 that this coupling can take place by the coincidence-controlled integration of a gene into the genome of a plant trans-
formed therewith. Prokaryotic genes, however, are combined
with preference in vitro with the suitable plant-specific
expression signals. A generally applicable method resides,
for example, in linking the 5' end of a structural gene
15 including the associated plant promoter with the coding sequence of the prokaryotic gene in such a way that a
fusion protein is synthesized in a plant cell transformed
therewith. This fusion protein consists of the N-terminal
amino acids of the plant protein and a predominant propor-
20 tion of the prokaryotic protein and, in its essential properties, corresponds to the prokaryotic protein
[R.T. Fraley et al., 1983, Expression of Bacterial Genes
in Plant Cells, Proc. Natl. Acad. Sci. USA 80 : 4803;
J. Paszkowski and M.W. Saul, 1986, Direct Gene Transfer
25 to Plants, in S.P. Colowick and N.A. Kaplan (ed.), Methods
in Enzymology 118 : 668]: A similar method utilizes solely
the plant promoter, including a portion of the 5'-untranslated
region of the transcribed m-RNA and makes do without the
5' end of the coding region of a plant gene. In the
30 expression in plants, an ATG codon of the prokaryotic
gene segment then serves as starting point for the trans-
lation. Analogously, the expression signals of plant
viruses can be employed [N. Brisson and T. Hohn, 1986,
Plant Virus Vectors; Cauliflower Mosaic Virus, in
35 S.P. Colowick and N.O. Kaplan (ed.), Methods in
Enzymology, 118 : 659]. Vectors for the expression of

116

prokaryotic genes in plants are known in the literature and are generally available.

A prerequisite for construction of a gene expressible in plants from a prokaryotic gene with the aid of the aforementioned vector plasmids resides in knowing the base sequence as present for the gene *tfdA*. Such knowledge permits exact localization of restriction cutting sites and thus makes it possible to link the various DNA segments in a base-accurate fashion. Furthermore, only such knowledge makes it feasible to perform controlled mutagenesis for the creation of novel, functionally significant DNA sequences, such as, for example, the introduction of novel restriction cutting sites.

Using this information, a plant/bacterial hybrid gene for 2,4-dichlorophenoxyacetic acid monooxygenase (DPAM) can be constructed and inserted into various plasmids in order to effect transfer and expression of the gene in plants. The flanking sequences to the bacterial gene, both 5' and 3', may be engineered by means known in the art for making the hybrid gene capable of expression in the plant. For example, constitutive or inducible promoters may be inserted at the 5'-end of the gene. In addition, 3'-signal sequences, such as, for example, termination and/or polyadenylation sequences, may be inserted in the plasmid to improve expression of the cloned gene.

The recombinant plasmid may then be transferred to plants by known methods; for example, either directly using vegetable protoplasts or via the Ti plasmid from *Agrobacterium tumefaciens* as described above. Other intervening steps known for tailoring plasmids for particular uses may be incorporated as well. If the Ti plasmid method is used, for example, intermediate plasmids may be constructed to allow expression of DPAM in a bacterial species, such as, for example, *E. coli*, which can carry and mobilize cointegrates of cloning vectors and disarmed Ti plasmids. *Agrobacterium* strains carrying the new plasmids are then used to infect plant cells.

17

Transgenic plants constructed according to these procedures can be grown in the presence of concentrations of certain substituted phenoxyacetic acids which are normally growth-inhibiting for untransformed plants. For example, tissue cultures of tobacco plants (*Nicotiana tabacum* W38) transformed with the DPAM gene of this invention under the control of a constitutive promoter from cauliflower mosaic virus demonstrate up to a 20-fold higher tolerance for 2,4-D than their untransformed counterparts (see Example 20) and transgenic plants grown in soil show at least a 10-fold higher tolerance.

This invention also includes the provision of antibodies, especially monoclonal antibodies, antibody fragments, for example, Fab fragments, and labelled antibodies which are capable of binding 2,4-D-monooxygenase, and which are prepared according to fully conventional techniques.

The invention will be explained in greater detail below with reference to illustrations and examples. In the examples, methods of genetic engineering have been utilized as they are generally known and well-tested in this field of art. Cloning of genes and analysis of gene structures were performed according to standard methods of genetic engineering as they have already been incorporated into the textbooks by E.-L. Winnacker ("Gene und Klon" [Genes and Clones], Chemie Publishers, Weinheim, 1985) and by R.W. Old and S.B. Primrose (Principles of Gene Manipulation, Blackwell Scientific Publ., Oxford, 1985). The methods and techniques have in each case been updated to the most recent status e.g. in I.K. Setlow and A. Hollaender (Genetic Engineering: Principles and Methods, Plenum Publ. Corp., N.Y.). Besides the specific publications, general working directions in the form of laboratory manuals have also been utilized in many instances [R.W. Davis et al., Advanced Bacterial Genetics: a Manual for Genetic Engineering; T. Maniatis et al., Molecular Cloning; T.J. Silhavy et al., Experiments with Gene Fusions, all from Cold Spring Harbor Lab., N.Y.;

A. Pühler and K.N. Timmis, Advanced Molecular Genetics, Springer, Berlin, 1984; D.M. Glover, DNA Cloning: a Practical Approach, JRL Press, Oxford, Washington, D.C., 1985].

- 5 On August 28, 1986, the following microorganisms were deposited with the Deutsche Sammlung von Mikroorganismen [German Collection of Microorganisms] (DSM) in Göttingen, Federal Republic of Germany (deposit number):

10	E. coli HMS 174 (pT7-5)	(DSM 3829)
	E. coli HMS 174 (pT7-6)	(DSM 3830)
	E. coli JA 221 (pGSS33)	(DSM 3831)
	E. coli LE 392 (pTJSS'035)	(DSM 3832)
	E. coli HB101 (pVK101)	(DSM 3833)
15	E. coli SK1592 (pKT231)	(DSM 3834)
	E. coli S17-1 (pKJS31)	(DSM 3835)
	E. coli S17-1 (pKJS32 RHAS')	(DSM 3836)
	E. coli S17-1 (pKJS(X)630)	(DSM 3837)
	E. coli SBC107 (pRME1)	(DSM 3838)
20	E. coli K38 (pGT1-2/pTJS'X535)	(DSM 3839)
	Alcaligenes eutrophus JMP134 (pJP4)	(DSM 3840)
	Alcaligenes eutrophus JMP222	(DSM 3841)
	Alcaligenes eutrophus JMP134:Tn5-2 (pJP4:Tn5-2)	(DSM 3842)
25	Alcaligenes eutrophus JMP134:Tn5-4 (pJP4:Tn5-4)	(DSM 3843)

Detailed Description of the Drawings

Abbreviations:

	Ap	- ampicillin resistance
	Km	- kanamycin resistance
5	kb	- kilobases
	M	- molecular weight marker
	kD	- kilodalton

Figure 1 - shows origin and restriction map of the
HindIII fragment cloned from pJP4, having a
size of 21 kb (Example 1) and of the thus-derived
subfragments (Examples 3-7), as well as the ex-
pression of phenoxyacetic acid degradation in
Alcaligenes eutrophus JMP222 by the correspond-
ing plasmids with a wide host range (Example 14).
The arrow marks the exact location of tfdA in
the plasmid pJP4.

Figures 2, 3, 4 - show the plasmids derived from pKT231
containing tfdA or parts of tfdA (Examples 3-7).
The thin lines denote the proportion stemming
from the vector plasmid; the fat lines denote
the insert stemming from pJP4.

Figures 5, 6, 7^a - show the plasmids derived from the
T7 promoter plasmids pT7-5 and pT7-6 with
tfdA-containing insert fragments (Examples 8-11).
The thin lines denote the proportion stemming
from the vector plasmid; the fat lines denote
the insert stemming from pJP4.

5-2-88

Figure 7b - shows the insertion of the omega fragment in the plasmid pTJS'X535 (Example 11).

5 Figure 8 - shows specifically radiolabeled proteins on the autoradiogram of a polyacrylamide gel, which proteins are expressed in high yield by the tfdA-containing plasmids with the aid of the T7 promoter (Example 15).

1: pTJSS'035; 2: pTJS'B435; 3: pTJS'X535;
4: pTJSS'036; 5: pTJS'B436; 6: pTJS'X536.

10 a denotes completely marked total cell protein,
b denotes protein specifically marked after induction of the T7 promoter.

15 Figure 9^{del 8} shows the shortened tfdA gene product expressed by plasmid pTJS'X535 omega with the aid of the T7 promoter (2b), as compared with the unshortened protein expressed by pTJS'X535 (3b) and with the insert-free vector plasmid pT7-5 (1b).

20 ~~* a denotes completely marked total cell protein,~~
~~* b denotes protein specifically marked after induction of the T7 promoter.~~

25 Figure 10 - shows the base sequence of the BamHI/SalI fragment having a length of 2058 bases, on which the gene tfdA is transcribed from the 5' end toward the 3' end. The arrows designate the beginning and end of the coding region of tfdA (Example 16).

FIGURE 11 Construction of plasmid pMCJC1007 for stable integration and expression of the DPAM gene in plants. In a first cloning step the procaryotic translation start codon GTG of gene *tfdA* was replaced by a synthetic oligonucleotide. To avoid dimerization at the BamHI site the oligonucleotide was linked in a two step ligation first to the BamHI end of linearized pUC18 and then to the DPAM coding sequence. In two following cloning steps the BamHI-SphI fragment containing the hybrid construct was inserted into the plant expression vector plasmid pA8, between the CaMV 35S promoter and the OCS 3' end. As pA8 contains a second SphI site outside of its polylinker, pA5 was used to form an intermediate plasmid based on pUC18.

FIGURE 12 Tolerance of transgenic tobacco against 2,4-D in tissue culture. Explants from *Nicotiana tabacum* W38 control plants (left) and transgenic plants (right) were incubated on medium containing 1.0 mg/l 2,4-D. Normal shoots develop from leaf discs which constitutively express DPAM (A, right). Transgenic explants with a light-inducible DPAM gene (B, right) show the same sensitivity against 2,4-D as the nontransformed tobacco W38 (left).

FIGURE 13 Cross tolerance of transgenic tobacco against other phenoxy herbicides. Substances listed below were incorporated into 2MS agar as described in the text. The graphic demonstrates the response of control tobacco (W38) and transgenic tobacco (WS7-7) against different synthetic auxins. The height of the bars indicates the concentration limit up to which cells from explanted leaf discs could differentiate into normal shoots. At higher concentration only callus growth was possible. 2,4-D: 2,4-dichlorophenoxyacetic acid, 4-CPA: 4-chlorophenoxyacetic acid, MCPA: 4-chloro-2-methylphenoxyacetic acid, 2,4,5-T: 2,4,5-trichlorophenoxyacetic acid, 4-CP: 2-(4-

chlorophenoxy)propionic acid, 2,4-DP: 2-(2,4-dichlorophenoxy)propionic acid, 2,4,5-TP: 2-(2,4,5-trichlorophenoxy)propionic acid.

S A and 14B
FIGURE 14 Effect of 2,4-D on transgenic tobacco plants. *S-2-a*

5 A, left: a *Nicotiana tabacum* W38 control plant; the plant on the right is a clonal replicate of the WS7-7 line, which expresses the DPAM gene under the control of the CaMV 35S promoter. Both plants were sprayed with 10 kg/ha 2,4-D-Na. B, left: a *Nicotiana tabacum* W38
10 control plant; the plant on the right is a clonal replicate of the WS4-7 line, which expresses the DPAM gene under the control of the light-inducible ST-LS1 promoter. Both plants were sprayed with 1 kg/ha 2,4-D-Na.

Example 1: Preparation of Plasmid pVJH21

(a) Isolation of Plasmids pJP4 and pVK101

Alcaligenes eutrophus JMP134 is incubated in 250 ml of PYF medium (peptone, 3 g/l; yeast extract, 3 g/l; fructose, 2 g/l) for 16 hours at 30° C. The plasmid pJP4 is isolated from the centrifuged cells in accordance with the procedure from J. Bacteriol. 135 : 227 (1978). The plasmid pVK101 [V.C. Knauf et al., Plasmid 8 : 45 (1982)] is isolated from E. coli HB101 according to the generally known process of alkaline lysis by T. Maniatis et al. (Molecular Cloning: a Laboratory Manual, Cold Spring Harbor Laboratory, Cold Spring Harbor, New York, 1982)

(b) Cloning of the 21-kilobase HindIII
Fragment from pJP4

A reaction mixture consisting of 20 µl (500 ng) of pJP4, 2.5 µl of TAS buffer (0.33 M tris-acetate, 0.65 M potassium acetate, 0.1 M magnesium acetate, 5 mM dithiothreitol (DTT), 30 mM spermidine, pH 7.9) and 2.5 µl (1 unit) of a diluted HindIII solution is incubated at 37° C for 120 minutes. A further reaction mixture consisting of 50 µl (14 µg) of pVK101, 5.5 µl of TAS buffer and 1 µl (20 units) of undiluted HindIII solution is incubated at 37° C for 90 minutes. Then 1 µl (30 units) of calf intestinal alkaline phosphatase (DIP, Boehringer, Mannheim) is added and the mixture is incubated for another 60 minutes. Respectively 9 µl of each of the two reaction mixtures is separated on a 0.7% low-melting agarose gel by electrophoresis. The gel is subsequently stained for 15 minutes in an ethidium bromide solution (5 µg/ml) and the DNA bands are made visible in UV 300 nm.

The individual 20 kb band of pVK101 restriction, as well as the second-largest band (21 kb) of pJP4 restriction are excised from the gel, both bands are combined, and the DNA is isolated from the agarose according to the procedure by T. Maniatis et al. (Molecular Cloning: a Laboratory Manual, Cold Spring Harbor Laboratory, Cold Spring Harbor, New York, 1982). The ethanol-precipitated DNA is taken up in 10 μ l of ligation buffer (66 mM Tris-HCl, 6.6 mM $MgCl_2$, 10 mM DTT, 1 mM ATP, pH 7.5), combined with 1 μ l (0.1 unit) of diluted T4-DNA ligase, and incubated for 16 hours at 14° C (= ligation batch). Competent cells of E. coli S17-1 [Biotechnology 1 : 784 (1983)] are produced according to the method by T.J. Silhavy et al. (Experiments with Gene Fusions, Cold Spring Harbor Laboratory, Cold Spring Harbor, New York, 1984), and 0.2 ml of the thus-obtained competent cells are mixed with 5 μ l of ligation batch, left for 40 minutes on ice, then heated for 3 minutes to 42° C (= transformation), subsequently diluted with 2 ml of LB medium according to T. Maniatis et al. (cf, above), and incubated for 60 minutes at 30° C (= phenotypic expression). Then portions of respectively 0.2 ml are spread out on LB agar plates containing 10 μ l/ml of tetracycline. The plates are cultured at 37° C for 16 hours. Several tetracycline-resistant colonies are found.

(c) Identification of pVJH2

In accordance with the process of alkaline lysis, an instant preparation of the plasmid DNA is performed on the thus-obtained tetracycline-resistant colonies, as described by T. Maniatis et al. The aforementioned process of treatment with alkaline phosphatase ensures that each of the tetracycline-resistant clones contains the desired insert. The recombinant plasmids are identified by restriction

enzyme digestion with EcoRI and subsequent electrophoretic separation of the DNA fragments in a 0.7% agarose gel according to T. Maniatis et al. (compare above). By comparing the sizes of the thus-obtained
5 EcoRI fragments with the size relationships known from the literature with respect to EcoRI and HindIII for pJP4 and pVK101 [R.H. Don, J. Bacteriol. 161 : 466 (1985) and V.C. Knauf et al., Plasmid 8 : 45 (1982)], an exact
10 identification is provided of the insert DNA as well as of its orientation in the vector plasmid pVK101.

(d) Restriction Map of the HindIII Insert
in pVJH21

A thus-obtained strain containing the plasmid pVJH21 is incubated in 400 ml of LB medium containing 20 µg/ml of tetracycline for 16 hours at 37° C.
15 From the centrifuged cells, the plasmid DNA is isolated according to the method of alkaline lysis by T. Maniatis et al. (see above). The isolated plasmid DNA of pVJH21 is digested in various batches with the enzymes EcoRI,
20 BamHI and SacI individually, as well as with the following nine combinations of several enzymes: EcoRI/BamHI (1); EcoRI/SacI (2); BamHI/SacI (3); HindIII/EcoRI (4); HindIII/BamHI (5); HindIII/SacI (6); HindIII/EcoRI/BamHI (7); HindIII/EcoRI/SacI (8); and HindIII/BamHI/SacI (9).
25 The DNA fragments are separated in a 0.7% agarose gel according to T. Maniatis et al. (see above), and their sizes are determined by comparison with HindIII-cut lambda-DNA. The restriction map is drawn with the thus-obtained data together with the size relationships known
30 for the plasmids pJP4 [R.H. Don, J. Bacteriol. 161 : 466 (1985)] and pVK101 [V.C. Knauf et al., Plasmid 8 : 45 (1982)] with respect to EcoRI, BamHI and HindIII.

Example 2: Production of Plasmid pGJS3

(a) Isolation of Plasmids pVJH21 and pGSS33

The recombinant strain E. coli S17-1 containing the plasmid pVJH21 prepared in Example 1, and
5 E. coli JA221 with the vector plasmid pGSS33 [G.S. Sharpe, Gene 29 : 93 (1984)] are incubated in respectively
300 ml of LB medium containing 20 µg/ml of tetracycline
for 16 hours at 37° C. The two plasmids are isolated from
the centrifuged cells in accordance with the method of
10 alkaline lysis by T. Maniatis et al. (see above).

(b) Cloning of SacI Fragments from pVJH21

A reaction mixture consisting of 10 µl
(5 µg) of pVJH21, 10 µl (1 µg) of pGSS33, 4 µl of TAS
buffer, 16 µl of water, and 0.5 µl (10 units) of SacI
15 is incubated at 37° C for 90 minutes. The restriction
enzyme is subsequently inactivated by heating to 68° C
for 15 minutes. The DNA is precipitated with ethanol
according to T. Maniatis et al. (see above), and the dried
precipitate is taken up in 40 µl of ligation buffer. After
20 addition of 1 µl (1 unit) of T4-DNA ligase, the ligation
batch is incubated for 16 hours at 14° C. E. coli LE392
[N.E. Murray et al., Mol. Gen. Genet. 150 : 53 (1977)] is
rendered competent for DNA reception according to the
method by T.J. Silhavy et al. (Experiments with Gene
25 Fusions, Cold Spring Harbor Laboratory, Cold Spring
Harbor, New York, 1984); 0.2 ml of competent cells is
combined with 5 µl of ligation batch, left on ice for
40 minutes, and transformed by 3 minutes of heating to
42° C. After dilution with 2 ml of LB medium and
30 incubation at 30° C, portions of 50 µl to 200 µl are
spread on LB agar plates containing 20 µg/ml of
chloramphenicol. The plates are incubated for 16 hours
at 37° C.

(c) Identification of Recombinant Plasmids

The thus-obtained chloramphenicol-resistant clones are spread parallel on two LB agar plates, one of which contains 25 µg/ml of streptomycin, the other of which contains 20 µg/ml of chloramphenicol. The plasmid DNA is isolated from clones that prove streptomycin-sensitive in this test with the aid of the instant method of alkaline lysis according to T. Maniatis et al. (see above). The size of the inserts contained in vector pVK101 is determined by restriction of the isolated plasmids with SacI, electrophoretic separation of the DNA fragments on a 0.7% agarose gel, and comparison of the bands visible in UV 300 nm with known fragments of a HindIII restriction of lambda phage DNA. The plasmid pGJS3 is a recombinant plasmid from pGSS33 and a DNA fragment having a size of 3 kilobases. Since the plasmid pVK101 per se has no scission site for SacI, the cloned DNA fragment clearly stems from the HindIII fragment, having a size of 21 kilobases, of pJP4.

Example 3: Preparation of Plasmids pKJS31 and pKJS32

(a) Isolation of Plasmids pGJS3 and pKT231

The strain E. coli LE392 produced in Example 2, containing the recombinant plasmid pGJS3, is cultured in 400 ml of LB medium with an addition of 20 µg/ml of chloramphenicol for 16 hours at 37° C. E. coli SK1592 with the vector plasmid pKT231 [M. Bagdasarian et al., Current Topics in Microbiology and Immunology, 96 : 47 (1982)] is incubated in 400 ml of LB medium with an addition of 50 µg/ml of kanamycin under the same conditions. The two plasmids are isolated from the centrifuged cells according to the process of alkaline lysis by T. Maniatis et al. (see above).

(b) Transcloning of the SacI Fragment
from pGJS3 into pKT231

A reaction mixture consisting of 40 µl (400 ng) of pGJS3, 4 µl (100 ng) of pKT231, 6 µl of
5 TAS buffer, 9 µl of water, and 1 µl (1 unit) of dilute
SacI solution is incubated at 37° C for 120 minutes.
The restriction enzymes are inactivated by subsequent
heating of the reaction mixture to 68° C for 15 minutes.
The batch is combined with 39 µl of water, 10 µl of 10-fold
10 concentrated ligation buffer, and 1 µl (0.1 unit) of a
dilute T4-DNA ligase, and the ligation batch is incubated
for 16 hours at 14° C. An amount of 0.2 ml of competent
cells of E. coli LE392, prepared according to the process
by T.J. Silhavy et al. (Experiments with Gene Fusion, Cold
15 Spring Harbor Laboratory, Cold Spring Harbor, New York,
1984) is mixed with 10 µl of the ligation batch, left
on ice for 40 minutes, heated for 3 minutes to 42° C,
mixed with 2 ml of LB medium, and incubated at 37° C for
60 minutes. Portions of 50 µl to 200 µl are spread on LB
20 agar plates containing 50 µg/ml of kanamycin. The plates
are incubated at 37° C for 16 hours.

(c) Identification of Recombinant Plasmids

The thus-obtained kanamycin-resistant
clones are spread in parallel on three different LB
25 agar plates, one of which contains 25 µg/ml of streptomycin,
the second containing 20 µg/ml of chloramphenicol, and
the third plate containing 50 µg/ml of kanamycin.
Clones that turn out in this test to be sensitive to
streptomycin and chloramphenicol, but resistant to
30 kanamycin contain, in the normal case, a recombinant
plasmid from the vector pKT231 (kanamycin resistance)
and from the 3-kilobase SacI fragment from pGJS3.
The size of the inserted DNA is determined by isolating

the plasmid DNA according to the conventional instant method by T. Maniatis et al. (see above), restriction of the DNA with SacI and subsequent analysis of the fragments by electrophoresis in a 0.7% agarose gel.

5 Plasmids having differing orientation of the insert in relation to the vector plasmid are identified by a restriction analysis with EcoRI carried out in the same way. The plasmid pKJS32 constitutes that form of the two possibilities wherein the EcoRI site of the insert
10 is located closest to the EcoRI site of the vector proportion. This plasmid yields two EcoRI fragments of a size of 0.9 and 15 kilobases. The plasmid pKJS31 represents the construction with the opposite orientation of the insert, yielding two EcoRI fragments of
15 the sizes 2.3 and 13.5 kilobases.

Example 4: Production of Plasmid pKJSB330

(a) Isolation of Plasmid pKJS31

The clone of E. coli LE392 obtained in Example 3 and containing the recombined plasmid pKJS31
20 is incubated in 400 ml of LB medium with an addition of 50 µg/ml of kanamycin at 37° C for 16 hours, and from the centrifuged cells the plasmid DNA is isolated by means of alkaline lysis according to T. Maniatis et al. (see above).

25 (b) Deletion of Smaller BamHI Fragment from pKJS31

A reaction mixture consisting of 10 µl (500 ng) of pKJS31, 7 µl of water, 2 µl of TAS buffer and 1 µl (1 unit) of a diluted BamHI solution is in-
30 cubated for 60 minutes at 37° C, and subsequently the enzymatic reaction is stopped by heating to 68° C for

15 minutes. The restriction batch is then combined with 50 μ l of water, 8 μ l of 10-fold concentrated ligation buffer, and 2 μ l (2 units) of T4-DNA ligase and incubated for 16 hours at 14° C. A mixture is prepared from 0.2 ml
5 of competent cells of E. coli S17-1, produced according to the conventional method by T.J. Silhavy et al. (Experiments with Gene Fusions, Cold Spring Harbor Laboratory, Cold Spring Harbor, New York, 1984), and 10 μ l of the ligation batch, left on ice for 40 minutes, heated to 42° C for
10 3 minutes, mixed with 2 ml of LB medium, and incubated for 60 minutes at 37° C. Portions of 50 μ l to 200 μ l are spread on LB agar plates containing 50 μ g/ml of kanamycin. The plates are incubated for 16 hours at 37° C.

(c) Identification of the Shortened Plasmids

15 The plasmid DNA is isolated from the thus-obtained kanamycin-resistant clones by means of the known instant method of alkaline lysis by T. Maniatis et al. (see above). By restriction of the plasmids with BglII and and electrophoretic separation of the DNA fragments
20 in a 0.7% agarose gel, plasmids are identified which have lost the small BamHI fragment from pKJS31. These plasmids possess only a single BglII scission site and therefore yield a BglII fragment having a size of 13.2 kilobases. They can thus be clearly differentiated from their origin
25 plasmid pKJS31 and are denoted as pKJSB330.

Example 5: Production of Plasmid pKJS32RHAS'

(a) Isolation of Plasmids pKJS32 and pRME1

The clone of E. coli LE392 obtained in Example 3 and containing the recombined plasmid pKJS32,
30 as well as E. coli SBC107 (pRME1) are cultured in respectively 400 ml of LB medium with an addition of

50 µg/ml of kanamycin at 37° C for 16 hours. The plasmids are isolated from the centrifuged cells by alkaline lysis according to T. Maniatis et al. (see above).

- 5 (b) Deletion of a HindIII/SalI Fragment
from pKJS32 and Subsequent Insertion of a
HindIII/SalI Fragment from pRME1

10 A reaction mixture consisting of 10 µl
(500 ng) of pKJS32, 10 µl (1 µg) of pRME1, 15 µl of water,
4 µl of TAS buffer, 0.5 µl (1 unit) of dilute HindIII
10 solution, and 0.5 µl (1 unit) of dilute SalI solution
is incubated for 120 minutes at 37° C, and the enzymatic
reaction is stopped by heating to 68° C for 15 minutes.
Then 10 µl of the restriction batch is combined with
25 µl of water, 4 µl of 10-fold concentrated ligation
15 buffer and 1 µl (1 unit) of T4-DNA ligase and the mixture
is incubated for 16 hours at 14° C. Cells of E. coli Sl7-1
rendered competent according to a conventional process by
T.J. Silhavy et al. (Experiments with Gene Fusions, Cold
Spring Harbor Laboratory, Cold Spring Harbor, New York,
20 1984) are mixed with 10 µl of the ligation batch, left on
ice for 40 minutes, heated for 3 minutes to 42°C, and, after mixing
with 2 ml of LB medium, incubated at 37° for 60 minutes. Portions of
5 µl to 200 µl are spread on LB agar plates containing 50 µg/ml of
kanamycin. The plates are incubated for 16 hours at 37°C.

- 25 (c) Identification of the Recombinant Plasmids

The plasmid DNA is isolated from the thus-
produced kanamycin-resistant clones by the conventional
instant method of alkaline lysis according to T. Maniatis
et al. (see above). By restriction with BamHI and
30 electrophoretic separation of the resultant DNA fragments,
plasmids are identified wherein, starting from pKJS32, the
smaller DNA segment between the HindIII cutting site located
on the vector portion in the middle of the kanamycin
resistance gene and the SalI cutting site located on the

insert has been removed and replaced by that HindIII/SalI
fragment from pRME1 carrying the 3' end of the coding
region of the kanamycin resistance gene. These plasmids
yield two BamHI fragments having the lengths of 13.3 and
5 2.0 kilobases and thus clearly differ from the starting
products and other possible byproducts of the ligation.
They represent structures wherein a 0.2-kilobase DNA segment
of the 3.0-kilobase insert of pKJS32 has been removed and
wherein a part of the kanamycin resistance gene has been
10 exchanged against another gene fragment of the same genetic
origin. These plasmids are called pKJS32RHAS'.

Example 6: Preparation of Plasmid pKJEΔB130

(a) Isolation of Plasmids pKT231 and pKJS32

The isolation of pKT231 was described in
15 Example 3, that of pKJS32 in Example 5.

(b) Cloning of the Smaller EcoRI/BamHI
Fragment from pKJS32 in pKT231

Two reaction mixtures, one consisting of
10 μl (500 ng) of pKJS32, 5 μl of water, 2 μl of TAS buffer,
20 and respectively 1 μl (1 unit) of dilute solutions of the
enzymes EcoRI, BamHI and PstI, the other consisting of
10 μl (250 ng) of pKT231, 6 μl of water, 2 μl of TAS buffer,
and respectively 1 μl (1 unit) of dilute solutions of the
enzymes EcoRI and BamHI, are incubated for 120 minutes at
25 37° C and, to stop the reaction, heated for 15 minutes to
68° C. Subsequently respectively 10 μl of the two restric-
tion batches are mixed, combined with 50 μl of water, 8 μl
of 10-fold concentrated ligation buffer and 2 μl (2 units)
of T4-DNA ligase, and incubated for 16 hours at 14° C.
30 A mixture is prepared from 0.2 ml of competent cells of
E. coli LE392, produced according to the process known in

the literature by T.J. Silhavy et al. (Experiments with Gene Fusions, Cold Spring Harbor Laboratory, Cold Spring Harbor, New York, 1984), and 10 µl of the ligation batch, left on ice for 40 minutes, heated to 42° C for 3 minutes, mixed with 2 ml of LB medium, and incubated at 37° C for 60 minutes. Portions of 50 µl to 200 µl are spread on LB agar plates containing 50 µg/ml of kanamycin. The plates are incubated at 37° C for 16 hours.

(c) Identification of Recombinant Plasmids

10 The thus-obtained kanamycin-resistant clones are spread in parallel on two LB agar plates, one of which contains 25 µg/ml of streptomycin, the other 50 µg/ml of kanamycin. The plasmid DNA is isolated from clones proving to be streptomycin-resistant in this test, in accordance with the instant method by T. Maniatis et al. (see above). Restriction of the DNA with XhoI and subsequent electrophoretic separation of the DNA fragments in a 0.7% agarose gel permit identification of recombinant plasmids yielding two XhoI fragments having the sizes of 3.0 and 9.7 kilobases.

20 When cutting the two plasmids simultaneously with EcoRI and BamHI, electrophoretic analysis yields two fragments having the sizes of 1.5 and 11.2 kilobases. The thus-identified plasmids are recombinants from the large EcoRI/BamHI fragment of pKT231 and the 1.5-kilobase EcoRI/BamHI fragment stemming from the insert DNA of plasmid pKJS32.

25 They are called pKJEΔB130.

Example 7: Production of Plasmid pKJS(X)630

(a) Isolation of Plasmid pKJSB330

The clone of E. coli Sl7-1, obtained in Example 4 and containing the recombinant plasmid pKJS330, is cultured in 400 ml of LB medium with an addition of 50 µg/ml of kanamycin at 37° C for 16 hours, and the plasmid DNA is isolated from the centrifuged cells by alkaline lysis according to T. Maniatis et al. (see above).

10 (b) Removal of a BamHI/XbaI Fragment from Plasmid pKJSB330

A reaction mixture consisting of 20 µl (1 µg) of pKJSB330, 15 µl of water, 4 µl of TAS buffer, 0.5 µl (2 units) of dilute XbaI solution, and 0.5 µl (2 units) of dilute BamHI solution is incubated at 37° C for 120 minutes. Then 20 µl of buffered phenol according to T. Maniatis et al. (see above) is added, the batch is mixed, then again blended after adding 20 µl of chloroform : isoamyl alcohol (24:1), and removed by centrifuging. The top aqueous phase is taken off and mixed with 120 µl of cold ethanol of a temperature of -20° C. The mixture is stored at -20° C for 30 minutes and then removed by centrifuging. The precipitated DNA is washed with cold 70% strength ethanol, dried under vacuum, and taken up in 20 µl of Klenow reaction solution [20 mM Tris-HCl, pH 8.0, 7 mM MgCl₂, 10 U/ml DNA polymerase I (= Klenow enzyme)]. After preliminary incubation at 37° C for 5 minutes, 2 µl of a solution of all four deoxynucleotide triphosphates (0.125 mM ATP, 0.125 mM CTP, 0.125 mM GTP, 0.125 mM TTP, Pharmacia) is added and incubation is continued for another 5 minutes. Then the batch is mixed with 80 µl of ligation buffer containing 25 U/ml of T4-DNA ligase and left at room temperature for 6 hours.

Then the mixture is incubated at 14° C for 16 hours,
0.2 ml of a suspension of competent cells of E. coli S17-1
prepared according to the method by T.J. Silhavy et al.
(Experiments with Gene Fusions, Cold Spring Harbor
5 Laboratory, Cold Spring Harbor, New York, 1984) is mixed
with 10 µl of the ligation batch, transformed, after 40 min-
utes of incubation on ice, by heat treatment (3 minutes,
42° C), mixed with 2 ml of LB medium, and incubated at
37° C for 60 minutes. Then portions of 50 to 200 µl of
10 the cell suspension are spread on LB agar plates contain-
ing 50 µg/ml of kanamycin. The plates are incubated at
37° C for 16 hours.

(c) Identification of the Shortened Plasmids

From the thus-obtained kanamycin-resistant
15 clones, the plasmid DNA is isolated by the conventional
instant method of alkaline lysis according to T. Maniatis
et al. (see above). By restriction of the plasmids with
SmaI and subsequent electrophoretic separation of the DNA
fragments in a 0.7% agarose gel, plasmids are identified
20 which have lost the DNA segment between the BamHI scission
site and the XbaI scission site on the insert of pKJSB330.
These plasmids yield, in the restriction analysis, two
SmaI fragments having a length of 2.6 and 9.9 kilobases
and are called pKJS(X)630. They are thus to be clearly
25 distinguished from the starting product pKJSB330.

Example 8: Production of Plasmids pTJSS'035 and pTJSS'036

(a) Isolation of Plasmids pT7-5, pT7-6 and pKJS32

Two strains of *E. coli* HMS174, one of which contains the plasmid pT7-5, the other the plasmid pT7-6, are cultured in respectively 400 ml of LB medium according to T. Maniatis et al. (see above) with an addition of 50 µg/ml of ampicillin at 37° C for 16 hours. From the centrifuged cells, the plasmid DNA is isolated by the known method of alkaline lysis by T. Maniatis. Isolation of pKJS32 is described in Example 5.

(b) Cloning of the *SacI*/*SalI* Fragments from pKJS32 in pT7-5 and pT7-6

A reaction mixture (A), consisting of 20 µl (1 µg) of pKJS32, 15 µl of water, 4 µl of TAS buffer, and respectively 0.5 µl (2 units) of the enzymes *SacI* and *SalI*, is incubated for 120 minutes at 37° C. Two further reaction mixtures, consisting either of 2 µl (1 µg) of pT7-5 (B) or 2 µl (1 µg) of pT7-6 (C), 15 µl of water, 2 µl of TAS buffer, and respectively 0.5 µl (2 units) of the enzymes *SacI* and *SalI*, are incubated under identical conditions. All three reactions are stopped by heating to 68° C for 15 minutes. Respectively 13 µl of the restriction batch (A) (pKJS32) is mixed with 7 µl of the restriction batch (B) (pT7-5) and, respectively, 7 µl of the restriction batch (C) (pT7-6). The two mixtures are combined with respectively 50 µl of water, 8 µl of 10-fold concentrated ligation buffer, and 2 µl (2 units) of T4-DNA ligase. Both ligation batches are incubated at 14° C for 16 hours. Respectively 0.2 ml of competent cells of *E. coli* LE392, prepared according to the procedure by T.J. Silhavy et al. (Experiments with Gene Fusions, Cold Spring Harbor Laboratory, Cold Spring Harbor, New

York, 1984), is mixed with respectively 10 µl of the ligation batches, left on ice for 40 minutes, heated to 42° C for 3 minutes, mixed with 2 ml of LB medium, and incubated at 37° C for 60 minutes. Portions of 50 µl to
5 200 µl are spread on LB agar plates containing 50 µg/ml of ampicillin. The plates are incubated at 37° C for 16 hours.

(c) Identification of Recombinant Plasmids

The plasmid DNA is isolated from the
10 resultant ampicillin-resistant clones by the known instant method of alkaline lysis according to T. Maniatis et al. (see above). By restriction of the plasmids with BglII and BamHI, as well as by dual restriction with BamHI and HindIII and subsequent electrophoretic separa-
15 tion of the fragments in a 0.7% agarose gel, plasmids are identified consisting of one of the two vector plasmids pT7-5 or pT7-6 and the small SacI/SalI fragment of pKJS32. These recombinant plasmids yield in this case, with pT7-5 being the starting material, two BglII
20 fragments having a size of 3.0 kilobases and 2.2 kilobases, a BamHI fragment of a size of 5.2 kilobases, and two BamHI/HindIII fragments having the sizes of 3.2 and 2.0 kilobases, and they are called pTJSS'035. In case of pT7-6 as the starting material, two BglII fragments
25 are obtained of a size of 2.8 kilobases and 2.2 kilobases, a BamHI fragment of a size of 5.0 kilobases, and two BamHI/HindIII fragments of the sizes 3.0 kilobases and 2.0 kilobases; they are called pTJSS'036. The re-combinant plasmids thus differ clearly from their
30 starting products pKJS32 and pT7-5 or pT7-6.

Example 9: Preparation of Plasmids pTJS'B435 and
pTJS'B436 :

(a) Isolation of Plasmids pT7-5, pT7-6
and pKJSB330

5 Isolation of pT7-5 and pT7-6 has been
described in Example 8; the isolation of pKJSB330 has
been disclosed in Example 7.

(b) Cloning of the Short SalI/BamHI Fragment
from pKJSB330 in pT7-5 and pT7-6

10 A reaction mixture (A), consisting of
40 µl (2 µg) of pKJSB330, 5 µl of TAS buffer, and re-
spectively 2.5 µl (4 units) of dilute solutions of the
enzymes SalI and BamHI, is incubated at 37° C for
120 minutes. Two further reaction mixtures, consisting
15 of 2 µl (1 µg) of pT7-5 (B) and, respectively, 2 µl (1 µg)
of pT7-6 (C), 20 µl of water, 3 µl of TAS buffer, and re-
spectively 2.5 µl (4 units) of dilute solutions of the
enzymes SalI and BamHI, are incubated under identical
conditions. The reactions are stopped by heating to
20 68° C for 15 minutes. The entire restriction batch (A)
(pKJSB330) is separated by electrophoresis in an agarose
gel [0.7% of agarose in TBE buffer according to
T. Maniatis et al. (see above) containing 0.5 µg/ml of
ethidium bromide, according to T. Maniatis et al. (see
25 above)]. Of the two DNA bands visible under UV light
(300 nm), the shorter band of 2.0 kilobases is isolated
from the gel by the method of DEAE membrane elution in the
following way: A suitably dimensioned section of a DEAE
membrane (Schleicher and Schuell, S & S NA-45) is
30 placed into a cut 1-2 mm below the band to be eluted.
The electrophoresis is continued until the respective

band has been entirely absorbed by the membrane. The membrane is then removed from the gel and washed for 10 minutes with 10 ml of NET buffer (150 mM NaCl, 0.1 mM EDTA, 20 mM Tris-HCl, pH 8.0). The washed
5 membrane is incubated in 150 μ l of HNET buffer (1 M NaCl, 0.1 mM EDTA, 20 mM Tris-HCl, pH 8.0) for 30 minutes at 68° C. The solution is removed and the membrane incubated another 10 minutes with 50 μ l of HNET buffer. The combined elution solutions (200 μ l) are diluted with 200 μ l
10 of water, mixed with 800 μ l of cold (-20° C) ethanol, and stored at -20° C for 30 minutes. The precipitated DNA is centrifuged and the precipitate taken up in 90 μ l of water. After adding 10 μ l of 3-molar sodium acetate solution according to T. Maniatis et al. (see
15 above), the mixture is reprecipitated with 200 μ l of cold ethanol (30 minutes, -20° C), centrifuged, the precipitate washed with cold 70% ethanol, dried under vacuum, and taken up in 80 μ l of ligation buffer. Of this mixture, respectively 40 μ l is combined with 3 μ l (100 ng) of
20 the restriction batch (B) (pT7-5) and, respectively, 3 μ l (100 ng) of the restriction batch (C) (pT7-6), and 2 μ l (1 unit) of T4-DNA ligase, and incubated at 14° C for 16 hours. An amount of 0.2 ml of competent cells of E. coli LE 392, prepared according to the method by
25 T.J. Silhavy et al. (Experiments with Gene Fusions, Cold Spring Harbor Laboratory, Cold Spring Harbor, New York, 1984), is mixed with 15 μ l of the ligation batch, left on ice for 40 minutes, heated to 42° C for 3 minutes, mixed with 2 ml of LB medium, and incubated at 37° C
30 for 60 minutes. Portions of 50 μ l to 200 μ l are spread on LB agar plates containing 50 μ g/ml of ampicillin. The plates are incubated at 37° C for 16 hours.

(c) Identification of Recombinant Plasmids

The DNA is isolated from the resultant ampicillin-resistant clones according to the known instant method of alkaline lysis by T. Maniatis et al. (see above). Restriction of the plasmids with EcoRI and subsequent electrophoretic separation of the fragments in a 0.7% agarose gel afford identification of plasmids containing the short SalI/BamHI fragment from pKJSB330 in one of the two vector plasmids. Recombinant plasmids with pT7-5 as the starting product yield two EcoRI fragments of 3.0 kilobases and 1.5 kilobases in size and are called pTJS'B435; those with pT7-6 as the starting material yield two EcoRI fragments of 2.8 kilobases and 1.5 kilobases in size and are called pTJS'B436.

Example 10: Production of Plasmids pTJS'X535 and pTJS'X536

(a) Isolation of Plasmids pT7-5, pT7-6 and pTJSS'035

E. coli LE392 containing the plasmid pTJSS'035 produced in Example 8 is cultured in 400 ml of LB medium with an addition of 50 µg/ml of ampicillin at 37° C for 16 hours. From the centrifuged cells, the plasmid DNA is isolated according to T. Maniatis et al. (see above). Isolation of pT7-5 and pT7-6 has been set forth in Example 8.

(b) Cloning of the Short SalI/XbaI Fragment from pTJSS'035 in pT7-5 and pT7-6

A reaction mixture (A), consisting of 6 µl (3 µg) of pTJSS'035, 15 µl of water, 3 µl of TAS buffer, and respectively 2 µl (5 units) of dilute solutions of the enzymes XbaI, SalI and BamHI, is incubated

at 37° C for 120 minutes. Two further reaction mixtures, consisting of either 2 µl (1 µg) of pT7-5 (B) or 2 µl (1 µg) of pT7-6 (C), 23 µl of water, 3 µl of TAS buffer, and respectively 1 µl (2.5 units) of dilute solutions of the enzymes XbaI and SalI, are incubated in the same way. The reactions are stopped by heating to 68° C for 15 minutes. In two different ligation batches, respectively 5 µl (500 ng) of the restriction batch (A) (pTJSS'035) is mixed with either 15 µl of restriction batch (B) (pT7-5) or 15 µl of restriction batch (C) (pT7-6), 50 µl of water, 8 µl of ligation buffer, and 2 µl (2 units) of T4-DNA ligase, and incubated at 14° C for 16 hours. A mixture is prepared from 0.2 ml of competent cells of E. coli LE392, prepared according to the method by T.J. Silhavy et al. (Experiments with Gene Fusions, Cold Spring Harbor Laboratory, Cold Spring Harbor, New York, 1984) and 15 µl of the ligation batch, left on ice for 40 minutes, heated to 42° C for 3 minutes, mixed with 2 ml of LB medium, and incubated at 37° C for 60 minutes. Portions of 50 µl to 200 µl are spread on LB agar plates containing 50 µg/ml of ampicillin. The plates are incubated at 37° C for 16 hours.

(c) Identification of Recombinant Plasmids

From the thus-obtained ampicillin-resistant clones, the plasmid DNA is isolated according to T. Maniatis et al. (see above). By restriction with the enzymes SmaI and EcoRI, as well as by dual restriction with EcoRI and SalI, plasmids are identified which contain the short, 1.4-kilobase SalI/XbaI fragment from pTJSS'035 in one of the two vector plasmids pT7-5 and, respectively, pT7-6. Recombinant plasmids with pT7-5 as the starting material yield two EcoRI fragments 3.0 and 0.8 kilobases in size, two SmaI fragments of the sizes 2.4 and 1.4 kilobases, and three EcoRI/SalI fragments of the sizes

2.4, 0.8 and 0.6 kilobases; they are called pTJS'X535. Recombinant plasmids with pT7-6 as the starting material yield two EcoRI fragments of the sizes of 2.8 and 0.8 kilobases, two SmaI fragments having the sizes of 2.2 and 1.4 kilobases, and three EcoRI/SalI fragments having the sizes 2.2, 0.8 and 0.6 kilobases; they are called pTJS'X536. The recombinant plasmids thus differ unequivocally from their starting materials.

Example 11: Preparation of Plasmid pTJS'X535omega

10 (a) Isolation of Plasmids pTJS'X535 and pDOC37

Two E. coli strains LE392, one of which contains the plasmid pTJS'X535 prepared in Example 10, the other the plasmid pDOC37, are cultured in respectively 400 ml of LB medium with an addition of 50 µg/ml of ampicillin at 37° C for 16 hours. The plasmid DNA is isolated from the centrifuged cells in accordance with T. Maniatis et al. (see above). The plasmid pDOC37 contains the omega fragment between two EcoRI restriction sites. Instead of pDOC37, it is likewise possible to utilize, as a source for the omega fragment, the plasmid pHP45omega described by Prentki and Krisch, 1984, in Gene 29 : 103.

(b) Cloning of the Omega Fragment from pDOC37 into the BglII Site of pTJS'X535

25 Two reaction mixtures, one of which consists of 2 µl (1 µg) of pTJS'X535, 14 µl of water, 2 µl of TAS buffer, and 2 µl (3 units) of BglII, the other of which consists of 2 µl (2 µg) of pDOC37, 14 µl of water, 2 µl of TAS buffer, and 2 µl (4 units) of a dilute BamHI solution, 30 are incubated at 37° C for 120 minutes. The reactions are stopped by heating to 68° C for 15 minutes. Respectively 10 µl of both restriction batches are combined and

mixed with 80 μ l of ligation buffer and 1 μ l (1 unit) of
T4-DNA ligase. The ligation batch is incubated at 14° C
for 16 hours. An amount of 0.2 ml of competent cells
of E. coli LE 392, prepared according to T.J. Silhavy et
5 al. (Experiments with Gene Fusions, Cold Spring Harbor
Laboratory, Cold Spring Harbor, New York, 1984), is mixed
with 10 μ l of the ligation batch, left on ice for 40 min-
utes, heated to 42° C for 3 minutes, mixed with 2 ml of
LB medium, and incubated at 37° C for 60 minutes. Portions
10 of 50 μ l to 200 μ l are spread on LB agar plates containing
50 μ g/ml of ampicillin and 80 μ g/ml of spectinomycin.
The plates are incubated at 37° C for 16 hours. By
selection with spectinomycin, it is ensured that all
growing clones contain a plasmid with the omega fragment
15 since the latter contains the gene for expression of
spectinomycin resistance.

(c) Identification of Recombinant Plasmids

The plasmid DNA is isolated from the
thus-obtained ampicillin- and spectinomycin-resistant
20 clones according to T. Maniatis et al. (see above). By
restriction with HindIII, the recombinant plasmids are
identified. If three HindIII fragments are thus obtained
having the sizes of 0.6, 2.0 and 3.2 kilobases, these in-
volve derivatives of pTJS'X535 which have the omega
25 fragment built in at the previous BglII scission site.
By ligation of the compatible overlapping ends of the
BglII and BamHI fragments, the restriction sites BglII and
BamHI at the linkage point are lost. These recombinant
plasmids are called pTJS'X535omega.

Example 12: Production of Phages MJSS'030 and MJSS'031

(a) Isolation of Plasmid pKJS32 and the
Double-Strand Forms of Phages M13tgl30
and M13tgl31

5 The strain *E. coli* JM101 [J. Messing et al.,
Nucl. Acids Res. 9 : 309 (1981)] as well as double-strand
DNA of the vectors M13tgl30 and M13tgl31 [M.P. Kieny et al.,
Gene 26 : 91 (1983)] can be obtained from Amersham Buchler
GmbH & Co. KG, Braunschweig. Double-strand DNA can also
10 be obtained according to the following method: 0.2 ml of
competent cells of *E. coli* JM101, produced according to
T.J. Silhavy et al. (Experiments with Gene Fusions, Cold
Spring Harbor Laboratory, Cold Spring Harbor, New York,
1984) are transformed with single-strand or double-strand
15 DNA of the M13 vectors according to a process known from
the literature, mixed with 3 ml of melted (45° C) H-Top
agar (10 g/l of tryptone, 8 g/l of NaCl, 8 g/l of agar),
and poured on H-agar plates (10 g/l of tryptone, 8 g/l of
NaCl, 12 g/l of agar). The cooled plates are incubated
20 at 37° C for 16 hours. 2 ml of TY medium (16 g/l of
tryptone, 10 g/l of yeast extract, 5 g/l of NaCl) is
inoculated with 20 µl of an overnight culture (16 hours,
37° C) of *E. coli* JM101 in TY medium and a single plaque
of the H-agar plate. After 16 hours of growth at 37° C,
25 the phage-infected cells, together with a fresh overnight
culture of *E. coli* JM101, are transferred by inoculation
into 400 ml of TY medium and cultured for 16 hours at
37° C. From the centrifuged cells, the double-strand
DNA is isolated according to T. Maniatis et al. (see
30 above). Isolation of plasmid pKJS32 has been described
in Example 5.

(b) Cloning of the SacI/SalI Fragment from pKJS32 into M13tg130 and M13tg131

A reaction mixture (A), consisting of 30 μ l (1.5 μ g) of pKJS32, 20 μ l of water, 6 μ l of TAS buffer, and respectively 2 μ l (4 units) of dilute solutions of the enzymes SacI and SalI, is incubated at 37° C for 120 minutes. Two further reaction mixtures, consisting of 10 μ l (300 ng) of double-strand DNA of M13tg130 (B) or M13tg131 (C), 25 μ l of water, 4 μ l of TAS buffer, and respectively 0.5 μ l (1 unit) of dilute solutions of the enzymes SacI and SalI, are incubated in the same way and heated, for stopping the reaction, to 68° C for 15 minutes. The entire restriction batch (A) is separated by electrophoresis in an agarose gel [0.7% agarose, 0.5 μ g/ml of ethidium bromide in TBE buffer according to T. Maniatis et al. (see above)]. Of the two bands visible under UV light, the smaller SacI/SalI band of 2.8 kilobases is isolated by means of a DEAE membrane from the gel in the same way as has been described for the 2.0-kilobase band in Example 9. The DNA is finally taken up in 80 μ l of water; of this amount, respectively 40 μ l is combined with 20 μ l of the restriction batches (A) (M13tg130) and, respectively, (B) (M13tg131) and 40 μ l of water, mixed with 100 μ l of buffered phenol/chloroform according to T. Maniatis et al. (see above), and centrifuged. The top aqueous phases are removed and, after adding 10 μ l of 3-molar sodium acetate solution according to T. Maniatis et al. (see above), mixed with respectively 200 μ l of cold (-20° C) ethanol. The mixtures are left at -20° C for 30 minutes, subsequently centrifuged, the precipitates are washed with cold 70% ethanol, dried under vacuum, and taken up in 40 μ l of ligation buffer. The two batches are combined with respectively 1 μ l (1 unit) of T4-DNA ligase and incubated at 14° C for 16 hours. Respectively 0.2 ml of

competent cells of E. coli JM101, prepared according to T.J. Silhavy et al. (Experiments with Gene Fusions, Cold Spring Harbor Laboratory, Cold Spring Harbor, New York, 1984), is mixed with 10 µl of the ligation batches, left
5 on ice for 40 minutes, heated to 42° C for 3 minutes, and combined with respectively 3 ml of melted (45° C) H-Top agar. The mixtures are combined with respectively 40 µl of IPTG (100 mM) and 40 µl of Xgal (2% in dimethylformamide) and poured on H-agar plates. The cooled plates
10 are incubated at 37° C for 16 hours.

(c) Identification of Recombinant Phages

Among the thus-obtained plaques, those stem from recombinant phages which do not exhibit a blue coloration. For further identification, the double-strand and
15 single-strand DNA is isolated from the colorless plaques according to methods known from the literature, compiled in "M13 Cloning and Sequencing Handbook" by Amersham. Restriction of the double-strand DNA with BglII and subsequent electrophoretic separation of the DNA fragments
20 identify derivatives of M13gt130 and, respectively, M13tg131 which contain the short SacI/SalI fragment from pKJS32. Recombinant phages with M13tg130 as the vector yield three BglII fragments of the sizes 0.7, 2.2 and 7.1 kilobases and are denoted as MJSS'030; those with
25 M13tg131 yield four BglII fragments of the sizes 0.6, 0.7, 2.2 and 6.6 kilobases and are called MJSS'031. They differ accordingly clearly from their starting products.

Example 13: Production of the tfdA Mutants *Alcaligenes*
eutrophus JMP134:Tn5-2 and JMP134:Tn5-4

(a) Transposon Mutagenesis of *Alcaligenes*
eutrophus JMP134

5 10 ml of PYE mdium (3 g/l of peptone,
3 g/l of yeast extract) is inoculated with 0.5 ml of an
overnight culture (16 hours, 30° C) of *Alcaligenes eutrophus*
JMP134 [R.H. Don et al., J. Bacteriol. 145 : 681 (1981)]
and shaken for 8 hours at 30° C. 10 ml of LB medium ac-
10 cording to T. Maniatis et al. (see above) is inoculated
with 0.1 ml of an overnight culture of *E. coli* S17-1
[Biotechnology 1 : 784 (1983)] containing the plasmid
pSUP2021, and shaken for 5 hours at 37° C. Subsequently,
the optical density at a wavelength of 600 nm is determined
15 in a photometer for both cultures. The cultures are mixed
in a proportion of 1 : 1 with respect to their optical
density and 1.5 ml of the mixture is distributed on a
PYE agar plate [R.H. Don et al., J. Bacteriol. 145 : 681
(1981)]. The plate is incubated for 16 hours at 30° C.
20 Subsequently the bacteria are rinsed off the plate with
1.5 ml of sterile 0.7% NaCl solution and the bacterial
suspension is diluted in two steps with 0.7% NaCl solu-
tion respectively 1 : 10 (0.1 ml + 0.9 ml). Of each
dilution, portions of 50 µl, 100 µl, 200 µl and 400 µl
25 are spread on minimal agar plates (1.6 g/l of dipotassium
hydrogen phosphate, 0.4 g/l of potassium dihydrogen
phosphate, 1 g/l of ammonium sulfate, 0.05 g/l of
magnesium sulfat x 7 water, 0.01 g/l of iron(II) sulfate x
7 water, 15 g/l of agar), these plates additionally con-
30 taining 2 g/l of fructose and 380 µg/ml of kanamycin.
The plates are incubated at 30° C for 3 to 7 days.

(b) Testing the Transposon-Containing
Strains for 2,4-D Degradation

The thus-obtained kanamycin-resistant clones represent progeny of the strain JMP134 which have integrated
5 the transposon Tn5 in a stable fashion in their genome. They are spread in parallel on two minimal agar plates, one of which contains 1 mM of 2,4-dichlorophenoxyacetic acid, sodium salt (2,4-D), the other of which contains 2 g/l of fructose and 380 µg/ml of kanamycin. The plates are
10 incubated for 3 days at 30° C. Strains exhibiting mutation in one of the genes responsible for 2,4-D degradation cannot utilize 2,4-D as a growth substrate (2,4-D-negative). They occur at a frequency of 0.1 - 1%, based on the entirety of transposon-containing, kanamycin-resistant clones.

15 (c) Identification of the Gene Defect of
the 2,4-D-Negative Mutants

The thus-obtained 2,4-D-negative mutants are spread in parallel on two minimal agar plates, one of which contains 2 mM of 3-chlorobenzoic acid, sodium salt
20 (3-CB), the other 2 g/l of fructose and 380 µg/ml of kanamycin. The plates are incubated for 3 days at 30° C. 2,4-D-negative transposon mutants of the strain JMP134 exhibiting mutation in the genes *tfdC*, *tfdD* and *tfdE* of 2,4-D degradation cannot utilize 3-CB as a growth sub-
25 strate (3-CB-negative), as could be shown by Don et al. [J. Bacteriol. 161 : 85 (1985)]. Mutants in the genes *tfdA* and *tfdB*, in contrast thereto, can grow on 3-CB as the sole carbon source (3-CB-positive). Transposon mutants
30 proving to be 3-CB-positive in this test are accordingly further investigated by means of an enzyme test for defects in the genes *tfdA* or *tfdB*. For this purpose, the strains are cultured in 250 ml of minimal medium (as above, without agar) with an addition of 15 mM of sodium pyruvate and 1 mM of 3-CB for 16 hours at 30° C.

The cells are harvested by centrifuging at 4° C, washed three times in 10 ml of cold (4° C) minimal medium (without carbon source), and recentrifuged, and finally suspended in such a quantity of minimal medium that an optical
5 density of 30 is obtained at 420 nm. For the enzyme test of 2,4-D-monooxygenase and, respectively, 2,4-dichloro-phenolhydroxylase, 1 volume proportion of the cell sus-
pension is mixed with 9 parts by volume of an oxygen-saturated minimal medium so that an optical density of 3
10 results at 420 nm. With the aid of a commercially available oxygen electrode, the oxygen absorption is then observed over 10 minutes without adding substrate. Sub-
sequently, 2,4-D is added to a final concentration of 1 mM, or 2,4-dichlorophenol is added to a final concentration of
15 0.2 mM, and the decrease in oxygen concentration over a period of 20 minutes is observed. By means of this test, tfdA mutants can be distinguished from all other types of mutations, as well as from the wild-type strain JMP134:
They show no increase whatever in oxygen consumption after
20 addition of 2,4-D, whereas, upon addition of 2,4-dichlorophenol, a significant rise in oxygen absorption occurs. The wild-type strain, as well as tfdB mutants, show increased oxygen consumption after addition of 2,4-D as the substrate and thus indicate intact 2,4-D-monooxygenase. The two
25 transposon mutants JMP134:Tn5-2 and JMP134:Tn5-4 are two 2,4-D-negative, 3-CB-positive mutants with a detectable 2,4-dichlorophenolhydroxylase, but without detectable 2,4-D-monooxygenase. They can thus be identified as tfdA mutants. This designation is confirmed by additional
30 experiments described in the examples below.

Example 14: Expression of Cloned tfdA Genes in
Gram-Negative Bacteria Other Than E. coli

(a) Preparation of Donor Strains for Conjugative
Transfer of tfdA-Containing Plasmids

5 The plasmids constructed on the basis of
mobilizable vectors with a wide hostrange, such as pVK101
[V.C. Knauf et al., Plasmid 8 :45 (1982)], pGSS33
[G.S. Sharpe, Gene 29 : 93 (1984)], and pKT321
[M. Bagdasarian et al., Current Topics in Microbiology
10 and Immunology 96 : 74 (1982)], the preparation of which
has been disclosed in Examples 1 through 7, can be trans-
ferred from the mobilizing strain E. coli S17-1
[Biotechnology 1 : 784 (1983)] by conjugation to other
gram-negative bacteria, such as, for example, Alcaligenes
15 eutrophus and Pseudomonas putida. For this purpose, they
must first be introduced by transformation into E. coli
S17-1 unless they have already been cloned in this strain.
To this end, the plasmid DNA is isolated from the clones
of E. coli LE392 containing the respective plasmid, in
20 accordance with T. Maniatis et al. (see above). A mixture
is made up of 0.2 ml of competent cells of E. coli LE392
produced according to T.J. Silhavy et al. (Experiments
with Gene Fusions, Cold Spring Harbor Laboratory, Cold
Spring Harbor, New York, 1984) and 1 µl of the isolated
25 plasmid DNA, left on ice for 40 minutes, heated to 42° C
for 3 minutes, mixed with 2 ml of LB medium, and in-
cubated at 37° C for 60 minutes. Portions of 50 µl to
200 µl are spread on LB agar plates containing the cor-
responding antibiotic suitable for selection. The plates
30 are incubated at 37° C for 16 hours.

(b) Conjugative Transfer

A thus-obtained strain of *E. coli* S17-7 containing one of the plasmids described in Examples 1-7 is cultured for 16 hours at 37° C in 5 ml of LB medium containing an antibiotic suitable for selection. As the recipient, one of the two *tfdA* mutants isolated in Example 13 (case A), the pJP4-free strain *Alcaligenes eutrophus* JMP222 [R.H. Don et al., J. Bacteriol. 145 : 681 (1981)] (case B), or *Pseudomonas* spec. B13 [E. Dorn et al., Arch. Microbiol. 99 : 61 (1974)] (case C) are cultured in 5 ml of PYE medium (3 g/l of peptone, 3 g/l of yeast extract) at 30° C for 16 hours. The cells of the donor strain are centrifuged and suspended in double the volume of PYE medium. The suspension of the donor cells and the culture of the recipient are mixed in equal parts. 100 µl of the mixture is pipetted onto a membrane filter (Millipore HA 0.45 µm) located on the surface of a PYE agar plate. Subsequently, the plate is incubated for 6 hours at 30° C. Then the cells are rinsed off the filter with 1 ml of 0.7% NaCl solution and the cell suspension is diluted in 7 steps respectively 1 : 10 (0.1 ml + 0.9 ml) with 0.7% NaCl solution. Of each dilution, 100 µl is spread on a minimal agar plate containing one of the following growth substrates: 1 mM 2,4-D in case A (*tfdA* mutant), 4 mM phenoxyacetic acid-Na in case B (*Alcaligenes eutrophus* JMP222) or 1 mM 4-chlorophenoxyacetic acid-Na in case C (*Pseudomonas* B13). The plates are incubated in case A for 14 days, in cases B and C for 4 days, at 30° C.

(c) Properties of the Newly Formed Strains

Mobilizable plasmids with a wide host range from the aforementioned group, containing an intact *tfdA* gene, render, after conjugative transfer, the respective
5 recipient strain capable of synthesizing a functional 2,4-D-monooxygenase. This leads in case A (*tfdA* mutant) to complementation of the mutation and thus to restoration of the wild-type property with respect to utilization of 2,4-D, i.e. strains are produced which are 2,4-D-positive,
10 growing on 2,4-D-minimal agar plates. The 2,4-D-monooxygenase coded by *tfdA* is furthermore capable of enzymatic conversion of, besides 2,4-D, also the compounds chemically related to 2,4-D, phenoxyacetic acid and 4-chlorophenoxyacetic acid, thus forming as the products phenol and, re-
15 spectively, 4-chlorophenol. Since the strain *Alcaligenes eutrophus* JMP222 does exhibit genes for the complete metabolizing of phenol, there result after conjugative transfer of *tfdA*-containing plasmids, in case B, strains having the novel property of utilizing phenoxyacetic acid
20 as growth substrate. In the same way, a *tfdA*-containing plasmid renders a *Pseudomonas* B13 equipped with this plasmid and having complete degradation pathways for phenol and 4-chlorophenol, capable of growing on the substrates phenoxyacetic acid and 4-chlorophenoxyacetic
25 acid (case C). The test for utilization of a specific substrate as described above is especially suitable for the confirmation of expression of the *tfdA* gene by DNA fragments cloned in mobilizable broad host range vectors, and the shortened derivatives of these fragments,
30 the production of which has been set forth in Examples 1-7. It turns out that the plasmids pVJH21, pGJS3, pKJS31, pKJS32, pKJSE330, pKJS32RHAS' and pKJS(X)630 all contain an intact *tfdA* gene whereas the plasmid pKJEΔB130 does not code an operable gene product.

Example 15: Accumulation of tfdA-Coded 2,4-D-Monooxygenase in E. coli

(a) Production of Highly Productive E. coli Strains

5 The recombinant plasmids described in Examples 8, 9, 10 and 11, produced based on the vectors pT7-5 and pT7-6, constructed by Tabor, contain cloned DNA fragments following a strong promoter of phage T7. Plasmids from this group, containing an intact tfdA gene
10 in the correct orientation with respect to the promoter, can produce 2,4-D-monooxygenase in large amounts under the influence of T7-RNA polymerase expressed by the plasmid pGP1-2 [S. Tabor et al., Proc. Natl. Acad. Sci. USA 82 : 1074 (1985)] in E. coli K38. Since the promoter
15 is recognized specifically only by T7-RNA polymerase, and since the latter is on the plasmid pGP1-2 under the control of a heat-sensitive lambda repressor, the synthesis of the gene product can be induced by heat. By adding rifampicin, it is furthermore possible to inhibit
20 the bacterial RNA polymerases. In order to produce the highly productive strains, the plasmid DNA is isolated from the plasmid-containing strains of E. coli LE392, obtained in Examples 8-11, according to T. Maniatis et al. (see above). E. coli K38, containing the plasmid pGP1-2,
25 is cultured for 16 hours at 30° C in LB medium that has been combined with 50 µg/ml of kanamycin for selection oriented toward pGP1-2. From the grown culture, competent cells are produced according to the method by T.J. Silhavy et al. (Experiments with Gene Fusions, Cold Spring Harbor Laboratory, Cold Spring Harbor, New York, 1984). An
30 amount of 0.2 ml of competent cells is mixed with 1 µl of the isolated DNA, left on ice for 40 minutes, heated to 37° C for 5 minutes, mixed with 2 ml of LB medium, and

incubated at 30° C for 60 minutes. Portions of 50 µl to 200 µl are spread on LB agar plates containing 50 µg/ml of ampicillin and 50 µg/ml of kanamycin. The plates are incubated at 30° C for 16 hours.

5 (b) Induced Production and Radiolabeling
of the tfdA Gene Product

The thus-obtained ampicillin- and kanamycin-resistant clones contain the plasmid pGP1-2 and one of the recombinant pT7 derivatives. They are
10 cultured in 10 ml of LB medium with an addition of 50 µg/ml of ampicillin and 50 µg/ml of kanamycin at 30° C. At an optical density of 0.5, measured at 590 nm, 0.2 ml of culture is removed by centrifuging. The cells are washed in 5 ml of M9 medium according to
15 T. Maniatis et al. (see above), again centrifuged, and finally suspended in 1 ml of M9 medium containing 20 µg/ml of thiamine and respectively 0.01% of all proteinogenic amino acids except for methionine and cysteine (18 in total). The cell suspension is shaken
20 at 30° C for 60 minutes; then the temperature is raised to 42° C. After 15 minutes, 10 µl of rifampicin solution (20 mg/ml in methanol) is added and the batch is left at 42° C for another 10 minutes. Thereafter the temperature is again lowered to 30° C; after 20 minutes the
25 batch is combined with 10 µCi of L-(³⁵S)methionine (Amersham, cell labeling grade) and left for 5 minutes at room temperature. The cells are subsequently removed by centrifuging, suspended in 120 µl of application buffer (60 mM of Tris-HCl, pH 6.8, 1% sodium lauryl
30 sulfate, 1% 2-mercaptoethanol, 10% glycerol, 0.01% bromophenol blue), and heated to 95° C for 3 minutes.

(c) Identification of Specifically Radiolabeled
tfdA Gene Product

The thus-obtained samples are loaded on
a 12.5% strength polyacrylamide gel [U.K. Laemmli,
5 Nature, 227 : 680 (1970)], and the entire bacterial
proteins are separated electrophoretically in accordance
with the method, known from the literature, described
therein. After electrophoresis has been completed, the
gel is stained for 30 minutes in a mixture of 2.5 g of
10 SERVA blue G, 454 ml of methanol, 92 ml of acetic acid,
and 454 ml of water and then decolorized for 24 hours in
a mixture of 50 ml of methanol, 75 ml of acetic acid, and
875 ml of water. The gel is dried on a filter paper
(Whatman 3MM) with the aid of a commercially available
15 gel desiccant; then autoradiography is performed according
to T. Maniatis et al. (see above) with an X-ray film
(Kodak Industrex AX), and the film is developed. On the
autoradiogram, an individual marked protein is identified
in those strains containing, besides pGP1-2, one of the
20 plasmids pTJSS'035, pTJS'B435, pTJS'X535 or pTJS'X535omega.
All of these homologous plasmids have the feature in
common that the cloned fragment is transcribed by the T7-
RNA polymerase toward the SalI end. In case of plasmids
having the reverse orientation of the insert (pTJSS'036,
25 pTJS'B435 and pTJS'X535), no specifically marked protein
is found. By comparison with standard proteins of a
known size, the molecular weight of the marked proteins
is determined. This results, for the gene products
expressed by pTJSS'035, pTJS'B435 and pTJS'X535, in a
30 molecular weight of 32,000, for the gene product
expressed by pTJS'X535 in a molecular weight of 29,000.

(d) Induced Production with Confirmation
of Enzymatic Activity

The above-obtained strains of *E. coli* K38, containing pGP1-2 and one of the plasmids pTJSS'035, pTJS'B435, pTJS'X535 or pTJS'X535omega, are grown at 30° C in 20 ml of LB medium with an addition of 50 µg/ml of ampicillin and 50 µg/ml of kanamycin. At an optical density of 1.0 at 590 nm, the temperature is increased to 42° C, and the culture is shaken for 25 minutes. Thereafter, 100 µl of rifampicin solution (20 mg/ml in methanol) is added and the culture shaken for 2 hours at 37° C. Detection of enzymatic activity of 2,4-D-monooxygenase in *E. coli* is carried out along the lines of the method of radioactive enzyme test described by Amy et al. [Appl. Env. Microbiol. 49 : 1237 (1985)], with the following modifications: The cells of an induced 20 ml culture are harvested by centrifuging, washed with 10 ml of EM medium, again centrifuged, and finally suspended in 10 ml of EM medium. The cell suspension is mixed in a 250 ml Warburg flask with 10 ml of transformation buffer, combined with 200 µg of unlabeled 2,4-D and 0.1 µCi of 2,4-dichlorophenoxy(2-¹⁴C)acetic acid (Amersham, 55 mCi/mmol), and shaken in the sealed flask at 21° C for 4 hours. Subsequently, 0.5 ml of beta-phenylethylamine is pipetted into the central vessel of the Warburg flask, and 2 ml of 1-molar sulfuric acid is injected into the cell suspension. After one hour of shaking at 21° C, the beta-phenylethylamine is taken up in 5 ml of counter fluid ("rotiszint 22", Roth), and the radioactivity of the sample is measured in a scintillation counter. Strains of *E. coli* K38 containing, besides pGP1-2, one of the plasmids pTJSS'035, pTJS'B435 or pTJS'X535 exhibit high enzyme activity in this test, whereas those with the plasmid pTJS'X535omega express no enzyme activity.

Example 16: Determination of Base Sequence of DNA
Cloned in M13

(a) Production of Deleted Phages from
MJSS'030 and MJSS'031

5 For sequencing the 2-kilobase BamHI/SalI
fragment of the recombinant phages according to the method
by Sanger [Proc. Natl. Acad. Sci. USA, 74 : 5463 (1977)],
a series of differently shortened phages is first produced
by controlled deletion according to a process known from
10 the literature [G. Henikoff, Gene, 28 : 351 (1984)],
making it possible to determine the entire base sequence
by overlapping sequencing of respectively 200 to 300 bases.
For this purpose, the double-stranded DNA is isolated
from strains of E. coli JM101 containing the phages
15 MJSS'030 and, respectively, MJSS'031 produced in Example 12,
in accordance with the method described therein for the
phage vectors M13tg130 and M13tg131. 10 µg of double-
stranded DNA of the phage MJSS'030 is cut with the
enzymes PstI and SalI, 10 µg of double-stranded DNA of
20 the phage MJSS'031 is cut with the enzymes BamHI and
SacI. The restriction batches are extracted with phenol
and the cut DNA is precipitated with ethanol. The subse-
quent digestion of DNA with exonuclease III and S1-nuclease,
Klenow capping of the projecting DNA ends, and ligation
25 take place exactly according to the method described by
Henikoff, with the following change: Instead of using
S1-nuclease, mung bean nuclease (Pharmacia) is utilized.
Portions of 0.2 ml of competent cells of E. coli JM101
[J. Messing et al., Nucl. Acids Res. 9 : 309 (1981)],
30 prepared according to the process by T.J. Silhavy et al.
(Experiments with Gene Fusions, Cold Spring Harbor
Laboratory, Cold Spring Harbor, New York, 1984), are
combined with respectively 20 µl of the ligation batches,
left on ice for 40 minutes, heated to 42° C for 3 minutes,

mixed with respectively 3 ml of melted (45° C) H-Top agar (Example 12), and poured on H-agar plates (Example 12). The plates are incubated at 37° C for 16 hours.

5 (b) Identification of Deleted Phages

E. coli JM101 is cultured in 2 ml of TY medium (Example 12) at 37° C for 16 hours. Of this culture, 1 ml is diluted with 100 ml of TY medium, and respectively 2 ml of the diluted cell suspension is infected with one of the above-obtained plaques. The cultures are shaken at 37° C for 5 hours and then centrifuged. From the supernatants of centrifugation, the single-strand DNA is obtained in accordance with the procedure indicated by Amersham [M13 Cloning and Sequencing Handbook, 1984]. The double-strand DNA is isolated from the centrifuged cells in accordance with T. Maniatis et al. (see above). The double-strand DNA of the shortened phages evolved from MJSS'030 is digested with BglII, that of the phages evolved from MJSS'031 with EcoRI and SalI. By subsequent electrophoretic separation of the fragments on a 2% agarose gel according to T. Maniatis et al. (see above), the size of the respective deletion is determined.

(c) Sequencing of Single-Strand DNA

25 The base sequence of the above-obtained single-strand DNA of the shortened phages, as well as of the single-strand DNA of MJSS'030 and MJSS'031 produced in Example 12 is determined according to the method, known from the literature, of dideoxynucleotide sequencing [Proc. Natl. Acad. Sci. USA 74 : 5463 (1977)]. In this procedure, the directions given by Amersham [M13 Cloning and Sequencing Handbook, 1984] are followed. In order to separate the DNA fragments, a gel having a length of 55 cm with a thickness increasing from 0.1 mm to 0.4 mm is utilized. The identified DNA sequences of

the shortened phages are joined with the aid of a computer program [C. Queen et al., Nucl. Acids Res. 12 : 581 (1984)] at overlapping regions to a DNA sequence encompassing the entire segment between the BamHI
5 scission site and the SalI scission site of the insert cloned in M13tg130 and, respectively, M13tg131. The base sequence is confirmed by comparison of the two thus-obtained complementary DNA strands.

(d) Properties of Sequenced DNA

10 With the aid of the above-mentioned computer program, all properties of the DNA are determined which result directly from the base sequence. Among these, the most important that can be cited are: the position of the recognition sites for restriction
15 endonucleases, the location and length of the open reading frames as possible coding regions of a gene, the frequency of specific bases and their distribution, and the occurrence of certain functional DNA sequences. Analysis of the DNA sequence shows an open reading frame,
20 the location, length and transcription direction of which coincide with the properties of the tfdA gene, described in Examples 14 and 15. This frame begins at a GTG starting codon with the base No. 748 and ends upstream of a TAG stop codon with base No. 1608, and therefore
25 has a length of 861 bases, corresponding to the length of the tfdA-coded protein (Example 15). The insertion of transcription and translation stop signals into the sole BglII cutting site of the sequenced fragment, as represented by the cloning of the omega fragment in
30 pTJS'X535 (Example 11), shortens this open reading frame by calculation to 768 bases, which coincides with the expression of a shortened and enzymatically inactive gene product by the plasmid pTJS'X535omega (Example 15).

Example 17: Construction of plasmids for constitutive and light-induced DPAM expression in plants.

a) Synthesis of oligonucleotides

To create a plant/bacterial hybrid gene for 2,4-dichlorophenoxyacetate monooxygenase (DPAM) the prokaryotic translation initiation codon GTG of gene *tfdA* has to be replaced by an ATG codon. This is achieved by a synthetic double stranded oligonucleotide which is designed to fit a 5' protruding end generated by FokI 8 bp downstream from the translation initiation codon. The oligonucleotide contains a sequence reconstituting the original open reading frame for DPAM, a consensus sequence for plant translation start sites upstream from the ATG start codon (Lütcke) and a 5' protruding end compatible with the cloning site BamHI (the sequences are specified in Figure 11). Oligonucleotides are made by an automatic synthesizer, like an Applied Biosystems 380A equipment, and purified by common methods as HPLC or preparative polyacrylamide gel electrophoresis which are described in the Applied Biosystems User Bulletin, issue No. 13, 1984 (Applied Biosystems, Foster City, California).

b) Construction of a hybrid coding sequence

Pairs of complementary oligonucleotides, each consisting of 20 bases, are phosphorylated by polynucleotide kinase, annealed by shifting temperature from 60°C to 37°C, and then ligated with pUC18 (Norranders et al., Gene 26:101, 1983) that had been digested with BamHI and SphI. The linear product is separated from an excess of oligonucleotides by agarose gel electrophoresis using DEAE membrane NA45 (Schleicher & Schuell). pTJSX535 plasmid DNA is digested with XbaI and SphI and separated by agarose gel electrophoresis. The 1.2 kb fragment containing the DPAM coding sequence is isolated from the gel by electroelution, partially digested by FokI, and repurified by agarose gel electrophoresis. DNA fragments of around 1.1 kb

61

isolated from the gel are ligated with the linear pUC18/oligonucleotide-construction, thus linking the oligonucleotide with the truncated coding sequence of tfdA (Figure 11). Competent cells of E. coli TB1 are transformed with the recombinant DNA and clones are selected on LB-agar containing 50 µg/ml ampicillin. E. coli TB1 is a derivative of strain JM101 (Yanisch-Perron et al., Gene 33:103, 1985) with the genotype: F'(traD36, proAB, lacI, lacZ ΔM15)Δ(lac, pro), SupE, thi, recA, Srl::Tn10(Tc^R) (Bart Barrel, personal communication). Clones containing plasmid pUJC1001 are identified by restriction analysis of isolated plasmid DNA. The correct ligation of the oligonucleotide is verified by sequence analysis of CsCl-purified pUJC1001 DNA using the dideoxy method (Sanger et al., Proc. Natl. Acad. Sci. USA 74:5463, 1977) modified for plasmid DNA as a template (Chen and Seeburg, DNA 4:165, 1985).

c) Construction of plasmids for plant expression

To fuse the new hybrid coding sequence at its 3' end to a termination and polyadenylation signal, DNA of pUJC1001 is digested with BamHI and SphI to excise the insert and then ligated with a BamHI and SphI cleaved pA5. pA5 is a pUC18 derivative that contains a 200 bp fragment of the octopine synthetase gene 3' end (OCS-polyA) (Dhaese et al., EMBO J. 2:419, 1983). The resulting plasmid pUJC01003 is cloned in E. coli TB1 as described in section b, above.

To fuse the 5' end of the hybrid DPAM gene to the cauliflower mosaic virus (CMV) 35S promoter (Paszkowski et al., EMBO J. 3:2717, 1984), DNA of both pUJC01003 and pA8 is digested with BamHI and SphI, mixed and ligated. The vector pA8 is based on the intermediate vector pMPK110 (Peter Eckes, Dissertation, Universität Köln, 1985) which is able to form cointegrates with disarmed Ti-plasmids of A. tumefaciens such as pGV3850kan (Jones et al., EMBO J. 4:2411, 1985). Competent cells of E. coli TB1 are transformed with the ligated DNA and clones are selected on LB agar containing 25 µg/ml

streptomycin. The resulting plasmid, designated pMCJC1007 (Figure 11), is identified by restriction analysis.

A similar construction is made using the light-inducible promoter of gene ST-LS1 (Stockhaus et al., Proc. Natl. Acad. Sci. USA 84:7943, 1987) instead of the constitutive CMV 35S promoter to form a plasmid designated pMLJC1005 (Figure not shown).

Example 18: Expression of the DPAM gene in E. coli.

For the two intermediate constructs, pUJC1001 and pUJC01003 (Figure 11), the DNA sequence predicts a fusion protein between the short N-terminal end of the lacZ coding region of pUC18 and the new hybrid coding sequence of the DPAM gene. This allows the expression of a functional DPAM in E. coli TBl from the IacZ promoter. Cells of E. coli TBl containing pUJC01003 are induced by IPTG and activity of DPAM within these cells is measured by the radioisotopic 2,4-D assay of Example 15d. DPAM activity is found in induced cells. In a similar assay nonradioactive 2,4-D is used as a substrate and the reaction product is detected by gas chromatography (Perkins et al., Weed Science 35:12, 1987). Subsequent mass spectrometry identifies 2,4-dichlorophenol (2,4-DCP) as the reaction product of the cloned DPAM.

Example 19: Transformation of tobacco with chimeric DPAM genes.

Both constructs containing the hybrid DPAM genes in combination either with the constitutive (pMCJC1007) or with the light-inducible promoter (pMLJC1005) are mobilized from E. coli TBl into the Agrobacterium strain C58C1 (pGV3850kan) (Jones et al., EMBO J. 4:2411, 1985) by a triparental mating (Sanchez-Serrano et al., EMBO J. 6:303, 1987) using E. coli GJ23 (Van Haute et al., EMBO J. 2:411, 1983) as a helper strain. Agrobacteria are checked by southern hybridization to contain the DPAM gene and subsequently used for leaf disc infection of

5 Nicotiana tabacum W38. Transformation and regeneration
of tobacco plants follow established procedures (Sanchez-
Serrano et al., EMBO J. 6:303, 1987). Shoots are regen-
erated from infected leaf discs by selection on 100 µg/ml
kanamycin and screened for nopaline production. Trans-
formed shoots are rooted and then transferred to soil.
RNA is isolated from leafs and analyzed by agarose gel
electrophoresis and subsequent northern blot hybridiza-
tion ^{using the 1.1 kb Bam HI - Sph I fragment of pTJSX535 as}
10 ~~using the 1.1 kb Bam HI Sph I fragment of pTJSX535 as~~
labeled probe. Transformed plants synthesize a
specifically hybridizing transcript of 1.3 kb in size.

**Example 20: Expression of the DPAM gene in tissue
culture of tobacco.**

15 As 2,4-D is known to promote callus growth and to
suppress shoot regeneration from explanted tobacco leaf
discs in tissue culture when applied in concentrations
higher than 0.1 mg/l, transgenic plants can be tested for
their ability to tolerate elevated levels of 2,4-D on a
20 synthetic "shoot regeneration" medium. This medium
consists of MS (Murashige and Skoog, Physiol. Plant
15:473, 1962), 15 g/l agar, 20 g/l sucrose, 1.0 mg/l 6-
benzylaminopurine and different concentrations of 2,4-D,
ranging from 0.1 mg/l up to 4.0 mg/l. Leaf discs from
25 sterile tobacco plants are placed on the agar and
incubated in the light for 3 weeks. Leaf discs from
transformed plants where the DPAM gene is under the
control of the constitutive 35S promoter develop normal
shoots on a medium containing up to 2.0 mg/l 2,4-D
30 (Figure 12A), whereas leaf discs from untransformed
control plants (Nicotiana tabacum W38) do not show any
differentiation on a medium containing more than 0.1 mg/l
2,4-D. This indicates that expression of the DPAM gene
can result in up to 20-fold higher tolerance against 2,4-
35 D.

Example 21: Response to differently substituted phenoxyacetic acids.

DPAM is able to degrade differently substituted phenoxyacetic acid derivatives. To show how transgenic tobacco expressing DPAM responds to the auxin activity of these compounds, the substances listed in Figure 13 are incorporated in MS-agar together with 1.0 mg/l BAP as described in Example 20. Leaf discs of transgenic tobacco expressing DPAM under the control of the constitutive 35S promoter and tobacco W38 are placed on the media and shoots arising are counted after 18 to 21 days. Figure 13 shows that transgenic plants express a similar tolerance to 4-CPA as to 2,4-D, a slightly reduced tolerance to MCPA and no tolerance to the tested tri-substituted phenoxyacetic acids and all tested phenoxypropionic acids.

Example 22: Resistance of transgenic plants to the herbicidal activity of 2,4-D.

Transgenic plants are transferred to soil and grown in the greenhouse. No difference in growth should be seen between untransformed and transformed tobacco. Control tobacco plants and transgenic tobacco plants are sprayed with 4 ml of a 1% aqueous solution of 2,4-D, Na-salt. With an estimated ground surface of 400 cm² per plant this roughly corresponds to an applicated amount of 10 kg/ha. Doses equivalent to 3 kg/ha and 1 kg/ha 2,4-D-Na are achieved by appropriate dilution of the spray solution. Whereas 1 kg/ha totally inhibits growth of control plants, transgenic plants which express the DPAM gene under the control of the constitutive 35S promoter are resistant to up to 10 kg/ha 2,4-D-Na (Figure 14A). Transgenic plants which express the DPAM gene under light-induction are at least resistant to 1 kg/ha 2,4-D-Na (Figure 14B).

In the foregoing description and examples, all temperatures are set forth uncorrected in degrees Celsius and, unless otherwise indicated, all parts and percentages are by weight.

The preceding examples can be repeated with similar success by substituting the generically or specifically described reactants and/or operating conditions of this invention for those used in the preceding examples.

Without further elaboration, it is believed that one skilled in the art can, using the preceding description, utilize the present invention to its fullest extent. The foregoing preferred specific embodiments are, therefore, to be construed as merely illustrative and not limitative of the remainder of the disclosure in any way whatsoever.

The entire texts of all applications, patents and publications cited above, and of corresponding application P 36 29 890.5 filed August 29, 1986 in the Federal Republic of Germany, are hereby incorporated by reference.

From the foregoing description, one skilled in the art can easily ascertain the essential characteristics of the invention and, without departing from the spirit and scope thereof, can make various changes and modifications of the invention to adapt it to various usages and conditions.